

PROCEEDINGS
OF THE
ROYAL SOCIETY OF EDINBURGH.

VOL. IX.

1875-76.

No. 95.

NINETY-THIRD SESSION.

Monday, 6th March 1876.

SIR WILLIAM THOMSON, President, in the Chair.

The following Communications were read :—

1. The Annual Periods of Thunder (with Lightning), Lightning (only), Hail, and Snow, at Oxford. By Mr Buchan.

During the twenty-one years ending 1873, the maximum period of thunder with or without lightning, at Oxford, extended from about 9th April to the end of October, the middle of the period being the first week of July; the three highest days taken consecutively being those immediately preceding the summer solstice. During the five months from November to March, only thirteen cases had occurred during the period. Lightning, on the other hand, had its maximum period from May to November,—particularly during August, September, and October. The maximum period of *hail* was during the first six months of the year; whereas, during the second half of the year, very few cases occurred. The snow period extended from the middle of October to the middle of May,—most falling from December to March,—the absolutely highest month being March.

Thus, *thunder with lightning*, at Oxford, closely follows the sun, the middle of the period being only about ten days after the

summer solstice; *lightning* (only) has its maximum period during that time of the year when the humidity of the air is at its maximum; *hail* is most frequent during that period of the year when the temperature is rising, or when the vertical layers of the atmosphere is in most unstable equilibrium; and *snow* during the coldest months of the year, with this striking peculiarity, that the maximum period is not in the depth of winter, but in March, in the end of winter; immediately after which the curve abruptly falls.

The intimate connection of the thunderstorm with summer rainfall, and the important bearing of the whole four curves on climatology, was referred to.

2. Note on the Origin of Thunderstorms. By Prof. Tait.

This Note does not refer so much to those great thunderstorms which extend over hundreds of miles in each direction, as to those small local storms which are often seen of from two to five or six miles only in diameter.

It refers particularly to those which are seen, in summer and autumn, to pass down the Tay valley. They almost invariably come from the westward, and I am told each is almost always accompanied by a storm of similar dimensions passing eastwards down the valley of the Forth. So far as I can ascertain, they seem both to commence almost abruptly somewhere in the district about Ben Ledi and Ben Lomond.

Seen from St Andrews, which they frequently pass at a few miles distance to the northward, they usually appear to be in a state of rotation about a vertical axis. It is not very easy to judge of the relative distances of the various clouds, so as to ascertain the *sense* of the rotation; but, in one case which I observed carefully last autumn, the rotation appeared to be in the *positive* direction,—i.e., opposite to that of the hands of a watch whose face is turned upwards.

If this be generally the case, and if it should be found that the direction of rotation of the companion storm in the Forth valley is *negative*, it would seem that their common origin may be explained on the following very simple hypothesis, which has the

additional recommendation of easily accounting for certain other singular phenomena.

It is known from balloon ascents that, in general, the atmosphere is arranged in horizontal strata of considerable depth or thickness, alternately moist and dry,—temperature diminishing steadily with increase of height in the moist, and remaining nearly constant throughout the dry, strata. These strata have usually horizontal velocities, differing (sometimes considerably) both in magnitude and direction. Thus near the common boundary of two such strata, fluid friction will in general tend to produce vortex motion,—the vortex columns being at first nearly horizontal, with their ends at the boundary, which is a surface of discontinuity.

A complete investigation of the possible circumstances would show *four* quite different cases:—

Vortex formed in $\left\{ \begin{array}{c} \text{dry} \\ \text{moist} \end{array} \right\}$ air, with its ends turning $\left\{ \begin{array}{c} \text{down} \\ \text{up} \end{array} \right\}$
into a stratum of $\left\{ \begin{array}{c} \text{moist} \\ \text{dry} \end{array} \right\}$ air.

The half vortex-ring thus formed tends, so far as it can, to become semicircular. It may thus extend downwards to the earth or upwards into the higher regions of the atmosphere.

If it extend downwards nearly to the earth, the lower portion will soon be destroyed by friction, and we shall have a couple of vertical vortex columns, with their ends respectively in the surface of discontinuity, and on the ground. They will of course rotate in opposite directions about the vertical, and their mutual influence will tend to cause them to progress in directions parallel to one another, the motion of each being in the same direction as that of the rotatory motion of the side which it at the moment turns to the other. This is exactly the presumed case of the little storms in the Tay and Forth valleys above referred to; the south side of the Tay column (that turned towards the Forth), moving eastward about the axis, while the axis itself moves to the east.

This theory is evidently capable of at once explaining the apparently *sudden* occurrence of such storms (of which waterspouts must be looked upon as small but quickly rotating examples), when the lower atmosphere has for hours been in a dead calm.

The disturbance has, in fact, its origin *above* the lower stratum, and works its way downwards into it.

It is also competent to explain the production of similar rotating storms in the higher regions of the atmosphere—many miles above the earth's surface—and thus to account for that by no means small number of cases of so-called "summer-lightning," which obviously cannot be explained by the occurrence of an ordinary thunderstorm at such a distance as to be below the spectator's horizon.

I have already explained to the Society that a possible source of at least a large part of the electric charge of a thunder-cloud is the contact-electricity of water-vapour and air. Thus while the precipitation of the vapour develops heat, the water particles precipitated are strongly electrified. And the aggregation into one of a number of equal little drops all charged to the same potential may increase the potential in any ratio whatever. Thus the charge on each drop in a large cloud may become so great that the electricity is driven entirely to the particles at its surface. This is supplementary to, and does not interfere with, Sir W. Thomson's explanation of the process by which the vapour is condensed.

It is possible that taking place in greatly larger spaces of air, but to a much smaller extent in each cubic foot, this sudden production, and as sudden scattering in all directions, of considerable quantities of electricity, may account for some of the main phenomena of the Aurora.

3. An Application of Professor James Thomson's Integrator to harmonic Analyses of Meteorological, Tidal, and other Phenomena, and to the Integration of Differential Equations. By Sir W. Thomson.


A first rough Model of Professor J. Thomson's Integrator was shown.

4. Note on the Thermo-Electric Position of Cobalt. By Professor Tait.

5. On a Glass Digester in which to Heat Substances under Pressure. By Dr E. A. Letts.

The objections to the use of sealed tubes are known to every practical chemist, and are a serious drawback to their employment. The chief of these are the time expended in the manufacture of the tubes, the amount of skill in glass-blowing required, the danger experienced in opening them, and above all, the fact that only a small quantity of material can be heated at one operation. Moreover, the same tube can seldom be used for more than three or four experiments, as each time it is sealed up its neck must be drawn out, and its length thus considerably decreased. These disadvantages were especially felt by me whilst preparing bromacetic acid, which was required in considerable quantities, and where as many as half a dozen tubes of bromine and acetic acid had to be heated before 100 grammes of the acid could be obtained. To obviate these objections I have had an apparatus constructed, which consists of a cylinder of glass, the walls of which are about half an inch thick. Its length is fifteen inches, its external diameter three inches, and its capacity about 600 cubic centimetres. At one end it is drawn out to a tube, whose aperture is only about one-sixth of an inch in diameter, though its walls are as thick as the rest of the apparatus. Originally this tube was provided with a stopcock, but at Professor Brown's suggestion, I have substituted a glass plate, which is ground flat, and accurately adapted to the top of the tube.

In order to keep the glass plate pressing on the tube the whole apparatus is placed in a frame, consisting of three brass wires arranged symmetrically around the cylinder, and attached by means of nuts, below, to a brass ring, and above, to a brass plate, through which latter a screw passes, which, when turned, presses on a brass plate placed on the glass cap.



As any experiments with such an apparatus would be attended

with danger, were it necessary to be in its neighbourhood, it occurred to me that an automatic arrangement might be employed to give notice that the temperature had been reached to which it was intended to subject the digester.

For this purpose I employed a thermometer with a somewhat wide tube and large bulb. A platinum wire is sealed into the bulb, and touches the mercury, whilst a brass wire passes down the tube, and is held in position by a binding screw. The two wires are connected with an electric bell, the brass wire being so adjusted, that when a particular temperature is reached the mercury touches its end, and thus completes the circuit, and causes the bell to ring.

In order to test the digester, about 200 grammes of a mixture of two-parts bromine and three of acetic acid was placed in it, and after fixing it in its frame, the whole apparatus was immersed in an oil bath and heated to 150° C., the temperature at which reaction in this case takes place. The experiment was made in a cellar, and the bell placed in a room some distance off. The gas to heat the oil bath was led by a tube from another cellar, so that it could be regulated without going near the digester. In about an hour and a half the bell rung, and thereupon the gas was shut off; and on examining the digester next day, it was found that the reaction had taken place, and that only twelve grammes of product had been lost—a very inconsiderable quantity.

As the action of bromine on acetic acid is very sudden, and accompanied by the disengagement of a large volume of hydrobromic acid, the apparatus may be considered to have undergone a very severe test, and that its efficacy for all ordinary purposes is established.

Should the digester come into general use, it will certainly save chemists much time and labour.

The following Gentlemen were elected Fellows of the Society :—

Rev. FRANCIS LE GRIX WHITE, M.A.

JAMES DUNCAN, Esq., Benmore.

Rev. NORMAN MACLEOD.

J. S. FLEMING, Esq.

JAMES DOUGLAS H. DICKSON, M.A. Glasg., B.A. Camb.

Monday, 20th March 1876.

SIR WILLIAM THOMSON, President, in the Chair.

The following Communications were read:—

1. On the Connection between Cohesion, Elasticity, Dilatation, and Temperature. By Professor George Forbes.

(*Abstract.*)

At various times there have arisen supporters of one or other of two extreme hypothesis concerning the nature of what we define as force. These are the hypothesis of "action at a distance" and of "no action at a distance."

According to the latter hypothesis, the centre of gravity of no body, however large or however small, can be moved from a position of rest, nor can its motion be altered in direction or amount, except by direct collision with another portion of matter.

Starting from this supposition as a basis of argument, and without assuming anything further as to the manner in which the different physical forces are caused by collisions, it is possible to arrive at some very general theorems; and from these theorems conclusions may be drawn as the nature and connection of some of the physical forces, which are necessarily true if the hypothesis of no action at a distance be true.

The principal result of these theorems is the following:—Let a rod be chosen of any substance whose cohesion and elasticity do not vary enormously with the temperature. Let a be its expansion, in terms of its length, when the temperature is raised 1° . Let β be the compression of the same rod, in terms of its length, when a unit weight is supported at its summit. Let c be the number of these units which, when suspended by the rod, suffice to break it by sudden rupture. Let θ be the absolute temperature at which all these experiments are made. Then the theory leads us to the conclusion that

$$a = \frac{\beta c}{\theta}.$$

Only a few experiments have been made by which we can test

this law. But the following values are the most accurate, and tend to prove the truth of the law. The apparent discordance in the case of iron is in part due to the variations in the qualities of that metal in different specimens.

	$\frac{\beta c}{\theta}$	α
Gold,	·00001484	·00001358
Silver,	·00001796	·00001809
Copper,	·00001511	·00001481
Platinum,	·00001006	·00000851
Iron,	·00001573	·00001220

In calculating this table, the values of c from the experiments of M. Wersheim are used; those of α from the experiments of Mr Mathiessen (except iron); those of β from the experiments quoted by Prof. Balfour Stewart in his Text-Book; and the assumed temperature is 18° C., or 283° absolute temperature.

2. Notice of the Completion of the Works designed by Sir Charles A. Hartley, F.R.S.E., for the Improvement of the Danube. By David Stevenson, Esq., V.P.R.S.E.

In 1868 I presented to the Society, on behalf of Sir Charles A. Hartley, a memoir published by the European Commission of the Danube, on the improvement of that river, and at the same time gave a notice of the works designed by Sir Charles Hartley for effecting that important object. These works have now been completed, and Sir Charles Hartley has again asked me to present to the Society a second memoir published by the Commission, which brings the history of the works constructed under their charge down to the time of their completion in 1873.

In supplement of the notice formerly communicated, which referred to a work in progress, it may not be uninteresting, now that the work is completed, to state briefly what has been effected by this most important and successful example of hydraulic engineering.

The engineering problem to be solved by the European Commission was the removal of the bar which obstructed the Sulina mouth of the Danube, which, in 1856, had a varying depth of channel never exceeding 11 feet. The design of Sir Charles Hartley—the engineer to the Commission—consisted in piers so constructed as to confine the current of the river in its passage into the Black Sea. At the date of my last notice the north pier had been extended to the length of 4640 feet, and the south pier to 3000 feet, and a maximum depth of $17\frac{1}{2}$ feet instead of 11 feet had been obtained. I, however, suggested in that notice, that as the Danube must continue to bring down an enormous mass of detritus, so in course of time the works which had proved so successful must be extended; and it appears that this has been found necessary, as the south breakwater, completed in 1871, has been extended to 3457 feet in length, and even with this additional length it is, I think, not improbable that in the course of time still farther extension may be required, for the Sulina mouth of the Danube will still discharge the same amount of water, bearing with it the same amount of alluvial matter, estimated in high floods at about 70,000 tons in twenty-four hours, the deposit of which at the extremity of the piers will still have a tendency, though in deeper water, to form a bar.

The works have, however, proved most successful, and reflect the highest credit on Sir Charles Hartley, by whom they were designed and executed, and the following is a summary of the results that have been obtained.

The total length of piers executed is 8789 feet, at a cost of L.185,352, being L.21 per lineal foot, in an average depth of 14 feet at low water. The navigable depth of the channel over the bar has been increased from 11 feet in 1856, to 20 feet in 1873. In 1853, 2490 vessels, of 339,457 aggregate tonnage, left the port; in 1869 there were 2881 vessels, with a tonnage of 676,960. Thus, while the number of vessels increased only at the rate of 16 per cent., the tonnage, due to the greater draught, had been increased at the rate of 50 per cent., a good practical proof of the value of the improvements. The number of shipwrecks at the mouth of the Danube has also been greatly diminished.

Monday, 3d April 1876.

SIR WILLIAM THOMSON, President, in the Chair.

The following Communications were read:—

1. Chapters on the Mineralogy of Scotland. By Professor Heddle. Chapter I.—On the Rhombohedral Carbonates. Communicated by Professor Tait.

Professor Heddle read a paper on the "Rhombohedral Carbonates occurring in Scotland," the first of a series of Chapters intended to embrace the analytical results of an investigation of all unknown or insufficiently determined Scottish species.

In this paper many analyses of the carbonates were submitted; and the pseudomorphic changes taking place in these were referred to in a special manner.

2. On Thermo-Dynamic Motivity. By Sir W. Thomson.
3. On the Vortex Theory of Gases, of the Condensation of Gases on Solids, and of the Continuity between the Gaseous and Liquid State of Matter. By Sir W. Thomson.
4. On two new Laboratory Apparatus. By William Dittmar.

The object of this communication is to submit to the notice of the Society two little inventions of mine, which, whatever may be the degree of originality which they can claim, will, I venture to hope, prove useful additions to the catalogue of chemical-laboratory appliances. The one is a new form of the precision balance, which pretends to execute *exact* weighings with a hitherto unattained degree of rapidity; the other is a contrivance for maintaining a constant pressure in a supply of gas, and thus making it possible, with comparative facility, to keep, say an air-bath, for any length of time, at a constant temperature.

The new *balance* differs from the instrument in its customary

form only in two points, of which the more important is a modification of the centre of gravity "bob" arrangement, which enables one, at a moment's notice, to shift the centre of gravity of the instrument from a certain definite position, I., to a certain other (higher) position, II., matters being arranged so that in passing from I. to II., the sensibility, *i.e.* the deviation, corresponding to an overweight of, say 1 milligramme, is increased in an exactly pre-determined ratio, such as of 1 : 10, for instance. For this purpose the "bob" is made very light, so that the distance through which it has to travel in order to effect the desired change of sensibility is not too small, and, instead of to a screw as usual, is fixed by *mere friction* to a vertical triangular steel rod forming part of the needle. The other new feature in the balance is, that the rider-principle, besides being discounted in a slightly different manner from the customary one, is extended to the determination of differences of weight up to 100 (instead of 10) milligrammes.




Fig. 1.

The arrangement adopted is represented in the accompanying sketch, for the interpretation of which it is only necessary to say that $C(10)$ and $(10)O$ are equal to $C(O)_1$ and $(O)_1(10)_1$ respectively, and that both $O(10)$ and $O_1(10)_1$ are each divided into 10 equal parts, the former by *notches* filed into the beam, the latter by *marks*; and to add, that there are two riders, one weighing p centigrammes for the left arm, and another weighing p milligrammes

for the right arm, the balance being adjusted so that, when both riders are at their zero-points, it is in equilibrium, and p being chosen so, that, supposing the large rider to be shifted to the n mark, and the small one to the m mark, this virtually amounts to the addition of $10n+m$ milligrammes to the charge in the right pan.

There is no need of my explaining how the balance is meant to be used; I will rather avail myself of this opportunity for drawing the attention of readers interested in the subject to a few inferences from the theory of the balance, which, obvious as they are, have hitherto not been sufficiently appreciated by either the authors of our physical handbooks or by practical balance-makers.

I. Given a balance in which everything is constant except the distance s of the centre of gravity of the empty instrument from the axis of rotation, and it is easily shown that (for a constant charge) the deviation α of the needle for a given over-weight Δ , and consequently the "sensibility" $\alpha = \frac{a}{\Delta}$, is the greater the less

s . This, of course, is duly stated by all authors; but what is always forgotten to be pointed out are two things, viz.—1st, That the "sensibility" has nothing to do with the *inherent precision* of the instrument; and 2dly, That supposing the sensibility to be increased, all the other good qualities of the balance get *less*; we diminish the rate of vibration (this rate being proportional to

$\sqrt{\frac{1}{a}}$); we diminish the range of differences of weight determinable by the method of vibration; we diminish the relative constancy (in opposition to variations in the charge) of the sensibility and the time of vibration. Considering now,—

II. The case of a *balance to be constructed*, the arm-length l and weight w of the empty beam also become variables, related to each other, according to some equation like $w = \text{const. } l$, and (assuming each of the pans to bear a certain medium charge p) we have

$$\frac{t}{\sqrt{a}} = \text{const. } \sqrt{l} \sqrt{\text{const.} + \text{const. } l},$$

i.e., by diminishing l we can increase the sensibility without diminishing the rate of vibration (or *vice versâ*); but the other

disadvantages mentioned under I. must be taken into the bargain, and, besides, the inherent precision of the balance gets less.* To pass to an example: What we gain by substituting a 7-inch for a 14 inch beam is that, for the *most convenient t*, the sensibility becomes 2 to 4 times greater; but this advantage is secured without expense in good qualities by placing before the graduated limb a lens magnifying the excursions of the needle into 2 to 4 times their natural size. This is the theory of the "*short beams*," which have lately come so much into fashion.

To come back to my own balance, I must not forget to thank Messrs Becker Sons of Rotterdam for the readiness with which they have, at their own risk, tried to realise my ideas in an actual instrument, which, by the way, is now being exhibited at South Kensington. To increase the usefulness of the instrument, I have caused Messrs Becker to add to it a glass plunger, which is adjusted so that it displaces exactly 10 grammes of water at 15°, and which consequently enables one with great rapidity to determine the specific gravities of liquids by the method of immersion.

To pass now to the new *gas-governor*, its most essential part consists of a mercury-manometer (fig. 2), of which one limb, A, is




Fig. 2.

about 20 mms. wide, and stands vertical; while the other, C, is of the width of a thermometer tube, and is placed horizontally.

* For fuller explanations, see my article "Balance" in the "Encyclopædia Britannica."

The empty part of the wider limb communicates, through F, with the gas-supply, through G with the gas-lamp serving to heat the air-bath; and the quantity of mercury is adjusted so that, when the gas is at the lowest pressure which, in the course of the experiment, it is likely to assume, the mercurial index in C occupies a certain convenient position *a*.

The manometer is connected with a constant battery (the circuit of which includes an electro-magnetic arrangement for opening or shutting the gas-tap), in such a manner that, as soon as, through an increase of pressure in E, the index in C travels ever so little towards the right of *a*, the current is closed, and the gas cut off.

The following Gentlemen were elected Fellows of the Society:—

JOHN MACMILLAN, M.A.

JOHN GIBSON CAZENOVE, D.D.

The following Gentlemen were elected Honorary Fellows of the Society:—

1. *Foreign Honorary.*

CARL LUDWIG, Leipzig.

FERDINAND DE LESSEPS, Paris.

2. *British Honorary.*

HENRY JOHN STEPHEN SMITH, Oxford.

THOMAS HENRY HUXLEY, London.

THOMAS ROMNEY ROBINSON, D.D., Armagh.

Monday, 17th April 1876.

Professor FLEEMING JENKIN in the Chair.

The following Communications were read:—

1. On an Improved Form of Galvanic Battery. By J. Cook, Esq. Communicated by Professor Tait.

I wish to direct attention to a simple improvement on battery cells, whereby porous cells are dispensed with, and the inconveniences of gravity batteries avoided.

I may say it is a year or two since I first tried the plan.

It consists in first filling the outer glass cell one-third or one-half with fine *silver* sand, then pushing a ring of glass (which I cut from a common ale pint-bottle with a hot soldering bolt) down an inch or so into the sand. The zinc element forms a ring round the glass, and the copper lies as a plate on the sand within the glass. Its superiority to the gravity batteries, and to those, such as Sir William Thomson's, where the sand forms a dividing layer between the copper with its sulphate below and the zinc with its liquid above, will be obvious. I did not find the cupric sulphate solution to diffuse into the zinc division. It so readily admits of inspection that it would be infinitely preferable to the Meidinger and other plans.

2. On the Properties of the Perigon Versor. By G. J. P. Grieve, Esq. Communicated by Professor Kelland.

3. Descriptions of some new or imperfectly understood Forms of Palæozoic Corals. By H. Alleyne Nicholson, M.D., D.Sc., F.R.S.E., Professor of Natural History in the University of St Andrews, and James Thomson, F.G.S.

(Abstract).

In this communication the authors gave descriptions of several new or imperfectly understood forms of Palæozoic corals. After giving a general account of the method of investigation employed by them, the genus *Heliophyllum*, Hall, was discussed at length. The external structure of this genus is very peculiar, and it was shown that the genus is not by any means as nearly related to *Cyathophyllum* as has been generally believed. The new genus *Crepidophyllum* was proposed for a group of forms possessing the extraordinary and characteristic endothecal dissepiments of *Helio-phyllum*, but with the remarkable character that the central tabulate area of the corallum is shut in by a well-developed accessory wall or inner mural investment. Sometimes this secondary investment constitutes a complete circular sheath to the central tabulate area, and in this case all the primary septa become

directly connected with the outer surface of the cylindrical tube thus formed. More commonly, the secondary investment is open all down one side, and becomes directly continuous with two of the primary septa, thus constituting a horse-shoe shaped space, formed by the central tabulate area together with a wide fossula containing three short septa. It was shown that the fine coral described by Mr Billings under the name of *Diphyphyllum Archiaci* was truly a *Crepidophyllum*. It was further shown that two different forms, of very similar aspect, had been included by one of the authors under the name of *Heliophyllum sub-cæspitosum*. One of these forms, the typical one, is a *Crepidophyllum*, and will stand as *C. sub-cæspitosum*. The other is a *Heliophyllum*, and the authors described this under the name of *H. elegantulum*.

The name of *Thysanophyllum* was proposed for a genus of æstræiform corals from the Carboniferous rocks of Scotland. This genus is related to *Lonsdaleia* in the general form of the corallum, in the presence of an exterior vesicular zone of large-sized cells, and in the possession of septa, which have no connection with the outer wall. It differs from *Lonsdaleia*, however, in the fact that the columella, so conspicuous in the latter genus, is wholly wanting, and the central area of the visceral chamber is occupied by strong remote, transverse tabulæ. Two species of the genus were described, under the names of *Thysanophyllum orientale* and *T. minus*.

Finally, the genus *Lindströmia* was proposed for a group of small corals, in which the corallum is simple and conical, with an extremely deep calice. The septa are well developed, and meet in the centre of the visceral chamber, where they coalesce to a greater or less extent, and form a strong twisted pseudo-columella, which projects into the floor of the calice, and occupies a large portion of the entire visceral chamber. There are no tabulæ, but the septa are furnished with more or less strongly developed dissepiments, which, however, are remote, and do not give rise to any vesicular zone. The genus may, perhaps, be regarded as belonging to the *Aporosa*. The species *L. columnaris* was described from the Devonian rocks of North America, and it was mentioned that the authors were in possession of other forms of the genus, still undescribed, from the Carboniferous rocks of Scotland.

4. On a Stable and Flexible Arch. By Professor
Fleeming Jenkin.

Monday, 1st May 1876.

Professor KELLAND, Vice-President, in the Chair.

The Council have awarded the Keith Prize for the biennial period 1873-75, to Professor CRUM BROWN, for his Researches on the Sense of Rotation, and on the Anatomical Relations of the Semi-circular Canals of the Internal Ear.

The following Communication was read:—

Is the Gaelic Ossian a Translation from the English?

By Professor Blackie.

The recent revival by a distinguished Celtic scholar of the theory of Laing that Macpherson's Gaelic Ossian is a translation from the English, affords an opportunity of examining that question in a more strictly philological fashion than it has hitherto had the fortune to enjoy. Parts of the question were no doubt touched by Mackenzie in the Report of the Highland Society, published in 1805 by Graham in his dissertation on the authenticity of Ossian, by Dr Clerk of Kilmalie, the distinguished author of the new version of Ossian in the late splendid edition published at the expense of the Marquis of Bute; but systematically grappled with the question has never been. Having recently gone through the whole of the originals, I have made careful notes of whatever might tend to settle this question, and have come to the conclusion, in the face of the statement of Mr Campbell—whose authority, no doubt, is one of the highest on the subject, that the Gaelic is unquestionably the original. The tests by which a translator's hand seems clearly discoverable are the following five:—(1) In the English version, awkward, forced, and unidiomatic expressions frequently occur, which can be clearly traced to the influence of a Gaelic original. (2) In all poems of any antiquity handed down

in manuscripts, difficulties will occur arising from obsolete words, errors in transcription, confused connection, and other causes. In such cases it is a common practice with translators to skip the difficulty, gloss over the matter with some decent commonplace, and sometime to make positive blunders, which is not difficult for a philologer to expose. All these signs of a translator's hand are frequent in Macpherson's English, and would be more so had he not indulged in such a habit of skipping generally that it is difficult to say in certain cases decidedly that the skip was made because the writer of the English wished to shirk a difficulty. (3) It is a common practice with translators, when they find a passage a little obscure, to remove the obscurity by some manifest alteration of the phrase, or even by interpolating a line, or interlarding a commentary. This also occurs in Macpherson. (4) It is not always that a translator writes under the same vivid vision, or the same fervid inspiration as the original poet; and the consequence is that he will occasionally degrade poetry into prose, and specially fail to bring out that individuality of character in his word-painting which Ruskin has so triumphantly insisted on in the case of the sister art. The instances of failure to seize the most striking features of the original, and the substitution of generic for specific epithets, are frequent in Macpherson. (5) Most translators yie —sometimes, no doubt, wisely—to the temptation of improving on their originals; and Macpherson, from what we know of him, was the last man in the world to think of resisting such a temptation. How much of the Gaelic, as we now have it—that is, his clean copy of his own originals—was subjected to this process of beautification, as we may call it, no one can now tell, but I have traced in several instances departures from the simplicity of the original Gaelic, which can be explained most naturally on the supposition that they proceed from a translator who has yielded, without any just cause, to this flattering seduction. When the results obtained by the detailed application of these tests are combined with the amount of external evidence to be found in the Highland Society's report to the effect that Macpherson actually did translate from Gaelic originals, and was seen by various parties for weeks and months employed in the work of translation, a cumulative proof was produced that he was most anxious to see by what arguments Mr

Campbell could rebut. If that gentleman, to whom Celtic literature owes so much (and who in fact is the Wolf of the Ossianic question), or any Galician who thinks with him, shall succeed in leading a counterproof, I can only conclude that, considering the scrappy and fragmentary nature of some of the materials in Macpherson's hands, it might possibly have been the case that the translator filled up some of the gaps in his tale in English, with the intention that they might be done into Gaelic before publication by Strathmaskie, Captain Morrison, or some other of his Highland coadjutors; but that the English, as a whole, is a translation from the Gaelic, and not a translation of the best quality in many respects, may be accepted as one of the best ascertained facts in the whole range of philological investigation.

The following Gentlemen were elected Fellows of the Society:—

Professor M. FORSTER HEDDLE.

J. F. RODGER, S.S.C.

WILLIAM THOMSON, F.C.S., Manchester.

Monday, 15th May 1876.

SIR WILLIAM THOMSON, President, in the Chair.

The Keith Prize for the biennial period 1873-75, which has been awarded to Professor CRUM BROWN for his Researches on the Sense of Rotation, and on the Anatomical Relations of the Semi-circular Canals of the Internal Ear, was presented by the President.

The following Communications were read:—

1. Notes on Dredging in Madeira, by the Rev. Robert Boog Watson, B.A., F.R.S.E., F.G.S.

The difficulties in the way of dredging at Madeira are many and considerable. This has probably prevented any of this work having been done since the publication of Mr Macandrew's list of Mollusca, presented to the British Association in 1854. The author

having dredged for several years at Madeira, confirms Macandrew's generalisation of the Mediterranean character of the Mollusca—excludes 12 of Macandrew's named list as having crept in by mistake, and to the 115 remaining species identified by Macandrew as Madeiran has added 200 to 250 more, making nearly 400 in all, of which 80 or perhaps 100 are probably new. These he hopes soon to publish.

2. Note on a New Fossil Foot-Print from the Permian Sandstone of Dumfriesshire. By Patrick Dudgeon, Esq., F.R.S.E. (Plate I.)

What appears to be an entirely new foot-print has lately been found in the red sandstone of this district. I have seen many of the foot-prints from the various quarries in the neighbourhood, but have not before observed this one, nor is it like any figured in Sir William Jardine's splendid work on the "Ichthyology of Annandale." The foot-prints in question were found in a bed about 20 feet from the surface, at Locharbrigg's Quarry, three miles from Dumfries. They exhibit the usual large hind and smaller fore foot; the impression of the hind foot measures $1\cdot5 \times 2\cdot6$, the fore foot $2\cdot3 \times 1\cdot9$; the stride of the animal appears to have been about 10'. The impression of the hind foot does not interfere with that of the fore foot, as is the case with several of the foot-prints figured in Sir William Jardine's work, the interval between them being 2': the hind foot, therefore, must have been put down in the rear of the fore foot when the animal was walking. The impression of the foot shows five toes, the *thumb* being placed far back. The most characteristic features in these foot-prints are the well-developed claws, and the oblique position of the toes, *i.e.*, they are placed to *march* one behind the other. In almost all the foot-prints I have seen, where the toes can be made out, the middle one appears the most prominent; this foot-print is markedly distinct in this respect.

As yet I have only been able to obtain one good specimen of this foot-print—a hind foot; the rest of the slab on which the casts were impressed was unfortunately used for a paving stone in a cottage in the neighbourhood. I got it lifted; but the rough





usage it had been subjected to had greatly injured the impressions on it; they were, however, sufficiently distinct to enable me to give the above particulars.

The accompanying photograph is a good one of the hind foot in my possession, about half the size of the original. The posterior pad of the foot is not quite complete, and it, together with the pads of the toes, are somewhat broken.

I would propose for these foot-prints the provisional name of *Herpetichnus loxodactylus*, the oblique-toed *Herpetichnus*, with the following abbreviated character:—

GENUS HERPETICHNUS, *Jardine* ("Ichnology of Annandale," 1853, p. 14).

Herpetichnus loxodactylus, sp. nov.

Sp. chars.—Fore foot = $2'3 \times 1'9$; hind foot = $3'5 \times 1'6$; stride about 10'; impressions free; toes 5, oblique; thumb far back; claws well developed.

Locality and horizon.—Permian Sandstone, Locharbriggs Quarry, three miles from Dumfries.

P.S.—In the discussion which followed this paper, Professor Huxley stated that so far as he could judge from the photograph exhibited, the markings closely resembled a foot-print he had described some years before in a paper read before the Geological Society of London, "On the *Stagonolepis Robertsoni* (Agassiz) of the Elgin Sandstones; and on the recently discovered Foot-marks in the Sandstone of Cummingstone" ("Quart. Jour. Geol. Soc.," 1859, xv. p. 440). The resemblance of these Cummingstone foot-marks to the *Chelichnus* of the Dumfriesshire flags was noticed by Professor Huxley in the paper referred to.

3. On the Decennial Period in the Mean Amplitude of the Diurnal Oscillation and Disturbance of the Magnetic Needle and of the Sun-spot Area. By J. A. Broun, F.R.S.

(Abstract.)

The author, in presenting results relating to the decennial period derived from observations made at Trevandrum during twenty-two years, has sought a redetermination of the mean duration of that

period, as shown by preceding magnetical observations connected with his own. The relation of the frequency and area of sun-spots to the amplitude of the diurnal movements of the magnetic needle gives an increased value to this investigation.

Two very different results have been obtained;—one by Dr Lamont, showing a period of 10·4 years; the other, by Dr R. Wolf, gives $11\frac{1}{9}$ years. Dr Lamont's result depends on the assumption that three periods occurred between 1787 and 1818—an assumption which is opposed to the conclusions which have been deduced from the sun-spot, auroral, and magnetic observations for that interval. Dr Wolf's result has therefore been accepted very generally by many of the most eminent scientific men in England and on the Continent.

The author determines the epochs of maximum and minimum range of the diurnal oscillations of the magnetic needle by the more exact method, in which the mean for twelve months corresponding (at its middle point) to each month of the year is obtained. Commencing with the Trevandrum observations, from the present time, proceeding backwards to the earliest series, showing a maximum, that of Cassini (Paris 1784–1788.) The maximum at this time (1787·25) is confirmed nearly by Gilpin's observations (London, 1786–1806). The latter do not show the minimum in 1792 and maximum in 1797, which should satisfy Dr Lamont's assumption, and they are considered by him, like the observations of sun-spots at the time, as worthless for this investigation. Dr Wolf, on the other hand, finds support in both for a minimum in 1798.

It is concluded by the author, from an examination of Gilpin's observations, that a maximum really happened in 1797·7, but so little marked as to make it probable that any slight corresponding increase of sun-spots would not be noticed by the single, not very accurate, observer at the time. Evidence, however, of a slight maximum is also found in Professor Loomis's investigation for the frequency of the aurora borealis. As it is certain that another maximum occurred about 1804 to 1806, the author finds that Gilpin's observations, which agreed with Cassini's at the commencement of the series, showed in all probability the true magnetic variations afterwards.

It results from these investigations that the mean duration of the

period is 10.45 years; but that it appears to undergo a variation between 8 and $12\frac{1}{2}$ years in an interval of 42 years. The small maximum of 1797.7, if a true result, may be expected to repeat itself at some future time, a result which could not fail to aid in the search for the cause of these variations.

The author shows, that according to the long period of 42 years, a *maximum* should have happened in 1776; but that year Dr Wolf has concluded to be one of *minimum* sun-spot frequency. That 1776 was really a year of maximum is confirmed by the observations of Van Swinden, who, it is shown, appears to have been the first to obtain a variation due to the decennial period, and to have pointed out the appearance of a law: it is also confirmed by the observations of Cotté at Montmerency.

The ratio of the ranges of the diurnal variation in the years when it is a maximum to that in the years of minimum, is compared for different parts of the world, and found nearly the same in both hemispheres. It is also found that the law of the diurnal variation is the same in the year of maximum as in the year of minimum. The author concludes that the increase of the diurnal variation is not due to a different cause from that which produces the variation at the minimum, and that this cause acts when there are no sun-spots in the same way as, though with less intensity than, when the latter have their maximum frequency and area. The magnetic variations are therefore not due to the sun-spots; the latter appearing only when the common cause produces diurnal variations having at least two-thirds of the maximum amplitude.

The results derived from the sun-spot area are compared with those from the magnetic observations. While a general agreement is found in the decennial variations from year to year, it is evident that the attempt to calculate the amplitude of the diurnal variation from the sun-spot frequency (as has been done by Dr Wolf) must give results frequently deviating widely from the truth, as might be expected from the previous conclusion.

The decennial period of disturbance of magnetic declination at Trevandrum, deduced from hourly observations in the eleven years, 1854 to 1864, is next considered. It is found that the mean disturbance at each hour of the day shows the decennial period; but that the range of the mean value, from the minimum to the maximum

year, is different for each hour, while the maximum and minimum do not happen at exactly the same time for all hours of the day. Secondary maxima and minima are also shown, which vary in their epochs gradually from midnight to noon.

No clear law appears to connect the amount of the maximum disturbance for any hour with that of the minimum for the same hour in the 11 years; the ratio of the first to the second is least for the hours near noon, and greatest for those near midnight. It is found, however, that the maximum and minimum mean disturbance in the diurnal variation for each year, as well as in the decennial variation for each hour, are connected by the following relations:— D_m being the maximum and D_o the minimum disturbance.

$$\sqrt{D_m} - \sqrt{D_o} = \text{Constant.}$$

The *monthly* mean disturbance at Trevandrum in each of the years 1854 to 1864 is compared with the monthly mean sun-spot areas deduced by Messrs De La Rue, Stewart, and Lœwy, from Carrington's and the Kew Observations, with the following result:—The monthly mean disturbance in the years 1854–56 had a considerable value, and marked variations when there were few or no sun-spots. In 1857 to 1862 there are found several maxima and minima of disturbance and sun-spots which occur simultaneously. In some cases, and especially in June 1862, there is a well-marked sun-spot area maximum without any corresponding change of magnetic disturbance. The cause of the solar disturbance did not then extend its action to the earth at that time.

The author concludes with a notice of the hypotheses proposed to explain the decennial period of magnetic variations and of sun-spot frequency, as well as of the cause of sun-spots. It is pointed out that no theory of sun-spot formation can be accepted which does not explain their non- (or very rare) appearance every 10 or 11 years, and therefore the cause of the decennial period is bound up with this explanation. A planetary action which disturbs the equilibrium of the solar gases has been proposed; no other seems to present itself, and this the author believes will be found ultimately to be in question; and though he has not himself been able to find any satisfactory evidence in its favour, yet remarkable results have been obtained by Messrs De La Rue, Stewart, and Lœwy.

4. On the Parallel Roads of Lochaber. By David Milne Home, LL.D.

(Abstract.)

The author referred to the papers written on the subject, beginning with that by Dr Macculloch, in the year 1817; and he explained the various theories suggested.

He intimated his adoption of the Lake theory, and expressed his adherence to the view he took in the memoir read by him in this Society in the year 1846, that the blockages of the lakes had been effected by detrital matter.

In support of this view, he pointed out that all over this district of the Highlands there were immense beds of clay, sand, and gravel up to the tops of the hills, at even 2000 feet above the present level of the sea.

These deposits he considered to be undeniable proofs of the prevalence of the sea over this part of the earth's surface to a height of 3000 feet at least.

When the sea retired, so as to expose to atmospheric action the higher parts of the country, there would be depressions in the surface of the land, where lakes would be formed. These lakes would continue at high levels, till the streams issuing from them cut through the detritus. In some cases, the process of erosion would be so gradual, that the lakes would subside without producing any conspicuous beach-lines on the mountain sides. In other cases, the removal of the blockages would be on a large scale, owing to the looseness of the detritus; and if these removals were separated from one another by a considerable interval of time, beach-lines of a permanent character would be formed on the sides of the mountains enclosing the valleys.

The author referred to the existence in this district of the Highlands, now, of several lakes at high levels, which were kept up by detrital blockages. He instanced, in particular, Loch Earba, in the Lochaber district, at a height of about 1150 feet, which was kept up by such means, and on whose banks there was evidence that the lake had once stood 30 feet higher than at present. Near Kingussie there was Loch Gwynae, at a height of 1015 feet above the sea, on whose sides there were traces of five terraces, the highest of which is 132 feet above the present surface of the lake.

He referred to the ample means of cutting through and removing detrital matter possessed by streams and rivers, mentioning particularly the enormous cliffs of detritus cut through by the Rivers Spean and Spey.

He next proceeded to discuss the theories of other geologists.

With regard to the theory that the parallel roads were formed by the *sea*, he adduced arguments to show, that this view was impossible, inasmuch as the "roads" should in that case have all stood at the same level; whereas, in the different glens, they stood at different levels. Moreover, it had been found, that old river courses existed, by which the water in Glen Gluoy flowed into the water in Glen Roy, and that the water in Glen Roy flowed through Glen Glaster into Loch Laggan,—a state of things utterly fatal to the marine theory.

With regard to the blockage of the lakes having been formed by *ice*, the following objections were stated:—

1st, The improbability that some of the glens were filled with water, whilst others were filled with ice, the temperature of those glens being all much the same, in consequence of nearly equal altitudes above the sea.

2d, The impossibility of getting a glacier to come to the exact spot, where the lakes stopped, to form barriers several miles long, so solid and permanent in structure, as to prevent the escape of the water from lakes above 300 feet deep.

The author concluded by referring to the numerous examples in the Lochaber district, of boulders perched on tops of hills, and of rocks smoothed and striated. These phenomena had been ascribed by some geologists to the action of land-ice. But, coupling with these high-perched boulders, the occurrence of kaims or eskars on the sides of the hills (above the parallel roads), and therefore formed before the Lake period, the author was inclined to ascribe these phenomena to one agent—viz., a sea loaded with ice, when the land was submerged, and to a strong current in the sea, from the north-west, which swept over the submerged land, and through such valleys as Glen Spean and Glen Roy. The lakes, he referred to the period when the land was rising out of the sea. Their beaches were formed on the marine detritus;—which also for a time dammed back the lake waters.

Monday, 5th June 1876.

D. STEVENSON, Esq., C.E., Vice-President, in the Chair.

The following Communications were read:—

1. Physical Observations in Northern Asia. By
Professor G. Forbes.
2. On Parallel Motions. By the Rev. John Wilson, M.A.,
Bannockburn Academy. Communicated by Professor
Kelland.

It has been well said that the transmission of force is an "essential condition in machinery." It is no less true that directness in transference is important; that the fewer links in the chain binding driver and follower together, the less likely is the machine to be put out of gear. There is no question here as to the comparative values of the different modes of conveying motion from a prime mover,—rolling contact, sliding contact, wrapping connectors, or linkwork. Each has its own excellencies; each its special advantages; and one is to be preferred to another only according to the nature of the work to be done.

I. *Watt's Parallel Motion*.—The general problem is the "commutation of circular with rectilinear motion." The importance of the question began to be felt soon after the introduction of the steam-engine; and Watt, in 1784, patented an invention which not only had the credit of being the earliest, but up to recent times, the most reliable and accurate parallel motion in existence. This system was a great advance on the huge chains and arches which were affixed to the working beam of the engine for the purpose of obtaining the desired motion; and it has proved to be sufficiently accurate for all practical purposes. The construction is simple, consisting of three bars: two, rotating round fixed centres, and connected at their other extremities to the third bar. A point in this bar, either within or without the points of junction, accord-

ing as the centres are on opposite sides, or on one side of the connecting rod, moves in a straight line. The simple explanation of the underlying principle is that the curvature in one direction is modified by the curvature in an opposite direction, due to the motion of the other bar. In both the original and the modified form of this three-bar motion there is a divergence from the straight line, which though inappreciable for small angles, becomes somewhat more apparent for larger ones.

The determination of the parallel point is given in the formula

$$QF : FP :: AP : BQ$$

In the figures 1 and 2, AP and BQ are the arms; PQ the con-




Fig. 1.

Fig. 2.

necting link. Let the centres A and B be chosen so that the line mM shall bisect the versed sines of BQ and AP; the throw QR being equal to PT.

$$\begin{aligned} \text{QF} : \text{FP} &:: \text{Lm} : \text{MK} : \text{Lq} : \text{Kp} \\ &:: \text{BQ} (1 - \cos \theta) : \text{AP} (1 - \cos \varphi) \\ &:: \text{BQ} \cdot 2 \sin^2 \frac{\theta}{2} : \text{AP} \cdot 2 \sin^2 \frac{\varphi}{2} \\ &:: \text{BQ} \cdot \theta^2 : \text{AP} \cdot \varphi^2 \quad . \quad . \quad . \quad (1) \\ &:: \text{BQ} \cdot \text{AP}^2 : \text{AP} \cdot \text{BQ}^2 \quad . \quad . \quad . \quad (2) \\ &:: \text{AP} : \text{BQ} \end{aligned}$$

(1) For small angles, $\sin \frac{\theta}{2} = \frac{\theta}{2}$ nearly.

(2) Ultimately, $BQ \sin \theta = AP \sin \phi$, or, $BQ : AP :: \sin \phi : \sin \theta :: \phi : \theta$.

In connection with Watt's system we are led to consider the motion of the pantagraph, which has been a means of extending the parallel motion. Thus, while to the point F is attached the air-pump rod of an engine; by means of the pantagraph, whose property is that it describes similar curves on a smaller or larger scale, another point will trace another parallel line, and to it therefore may be fixed the end of the piston-rod. By superadding the parallelogram of bars to the Watt parallel motion, and making E,—the angle of the parallelogram—the connecting point of the piston rod, so as to concentrate the force; we have the point E describing a curve similar to F, when

$QF : BQ :: PQ$ or $HE : BH$. (Fig. 3.)




Fig. 3.

The problem of the pantagraph may be proved as follows. Let the initial position be at an angle of 90° , the arms lying along the axes of x and y . (Fig. 4.)

Given the fulcrum A and the ratio $AB : AC$, to find B_1 , the position which B the pencil assumes when the tracer C moves to C_1 . When the position of C_{11} is given, the angle C_1D_1A is known, for AD_1 and CD_1 are constants. Join BB_1 and CC_1 ; AB_1 and AC_1 (which latter are not assumed to be coincident).

In the triangles C_1D_1A and B_1F_1A , the angle B_1F_1A is equal to the angle C_1D_1A ; and the sides about these angles are propor-

tional, viz., $B_1F_1 : F_1A :: C_1D_1 : D_1A$. Therefore the triangles are equiangular (Euc. VI. : 6).

Wherefore the angle F_1AB_1 is equal to the angle D_1AC_1 , that is they are identical; and AB_1C_1 is a straight line. Now in the triangle ACC_1 , — $AB : AC :: AB_1 : AC_1$. Therefore BB_1 is parallel to CC_1 ; and $BB_1 : CC_1 :: AB : AC$, which is the given ratio.




Fig. 4.

Wherefore the position of B_1 is found. And any other position such as C_n will have its projection in similar ratio to the given one $AB : AC$ or $AF : AD$.

II. *The Reciprocator*.—About 1864 M. Peaucellier communi-

cated to the Société Philomathique a paper anent a newly-discovered means of producing parallel motion; but, at the time, little notice was taken of the subject. When Lipkin, a pupil of Tchebicheff, rediscovered it a few years later, it awakened the attention of the most eminent mathematicians of the day; and now it promises to become a power in the field of higher analytical investigation. The marvellous extension of the problem is due chiefly to the research of Professor Sylvester, who was the first in this country to direct attention to it.

The fact of pure linkwork constituted the difference between Peaucellier's and all other attempts at parallel motion. These have been only approximate, or if exact, the slot has been called into action; while in the case before us the motion is entirely due to the grouping of links around fixed centres.

The system consists of seven links. (Fig. 5.) A cell or rhombus $MPNC$ is jointed at two opposite angles to two links AM and AN , which are called connectors. To one of the remaining angles is attached a radial bar BP . Of the two fixed points, A is called the fulcrum; B , the pivot. The points C and P are the poles of the cell. The variable distances \overline{CA} and \overline{AP} are called the arms of the cell, while the difference between the squares of AM and MP is termed the modulus.

When A, the fulcrum, lies without the rhombus, the cell is called a positive cell; when it falls within, the cell is negative.

The following results can easily be verified :—

(1.) \overline{APC} is a straight line.

(2.) $\overline{AP} \cdot \overline{AC}$ is a constant quantity (Euclid III. 36), \bar{N} being at every moment the centre of a circle with a radius \overline{NP} or \overline{NC} .

This result is expressed

by saying that the curves described by P and C are the inverse of one another. Now the inverse of a circle is generally another circle. If, then, one of the poles of the cell be made to revolve in a circle round B, the other pole will describe




Fig. 5.

a circle. There is however an exception, for when the fulcrum is in the circumference of the circle described by P or C, the other pole describes a straight line.

The following is the geometrical proof:—

Let the positions of P be p_1, p_2, p_3 , &c., and the corresponding positions of C, c_1, c_2, c_3 , &c.; and suppose that the initial position of the system is when AC lies along the axis of x , that is, in a straight line with the centres A and B. Join P and p_1 ; C and c_1 . Join the corresponding pairs Pp_3, Cc_3 , &c.

By Eucl. II.: 12. $AM^2 = MP^2 + AP^2 + 2AP \cdot PK$

$$Am^2 = mp^2 + Ap^2 + 2Ap \cdot pk$$

$$\therefore AP [AP + 2PK] = Ap [Ap + 2pk]$$

$$AP \cdot AC = Ap \cdot Ac.$$

$$\text{or } \frac{AP}{Ap} = \frac{Ac}{AC}$$

In the triangles APp and ACc the angle A is common ; and the sides about it are proportional, viz.,

$$AP : Ap :: Ac : AC$$

Therefore the triangles are equiangular (Euc. VI. 6), wherefore the angle $c_1CA = Pp_1A$. But Pp_1A , the angle in a semicircle, is a right angle (Euc. III. 31), therefore c_1CA is also a right angle.

With any other position p_2 of P, C will take up a corresponding one c_2 , by joining which with C a right angle is formed. Wherefore the point C moves in a straight line at right angles to AB produced.

When the ratio subsisting between the length of the radial bar and the distance of the pivot from the fulcrum is not one of equality, the position of the parallel point is a circle, which is concave or convex with respect to the fulcrum, according as BP is greater or less than AB. The results, as determined by methods of analysis, show that the equation is the same for both the symmetrical and non-symmetrical forms of the cell; that is, for both the ordinary cell, where the pivot B in the initial position lies in a line with the fulcrum and the poles,—and for that form where the line passing the poles makes in the initial position a tangent to the circle described by the radial bar. Thus, if R were the centre of the circle described by C, its distance from the fulcrum A,

$$\text{is } RA = \frac{AB [AM^2 - MP^2]}{BP^2 - AB^2}; \text{ and the length of the radius}$$

$$\text{is } RC = \frac{BP [AM^2 - MP^2]}{BP^2 - AB^2}$$

The determination of the position of R, which is evident in the non-symmetrical form of the cell, is simply given in the ratio $RC : RA :: BP : AB$.

Thus, if $AM = 7$; $MP = 5$; $BP = 3$; and $AB = 1$

$$\text{then } RC = \frac{BP [AM^2 - MP^2]}{BP^2 - AB^2} = \frac{3(49 - 25)}{9 - 1} = 9$$

$$RA = \frac{AB}{BP} \cdot RC = 3$$

One form of the cell has given rise to a discussion of the question whether the parallel motion of Peaucellier is not simply a modification of the pantagraph. The resemblance between the two systems is not noticeable at first sight; and one would be inclined to deny any connection between them.

In this system the connectors are joined not at the opposite angles of the rhombus but at such points in the adjacent sides produced that at every moment they are parallel to the remaining

sides (fig. 6). APC

is a straight line.

$AM = CM$ since $AM \parallel LP$.

Let \overline{LP} be produced to

F, making $\overline{LF} = \overline{LM}$, and

complete the rhombus

LFKM. The bars \overline{AN} ,

\overline{CN} , and \overline{IP} can be re-

removed without interfer-

ing with the motion of

\bar{C} and \bar{P} . Thus AM,

MC, with LF, FK form

 \bar{C}, \bar{P} takes another, equi

III. *The Gorgon Line*

Russell is exact, and co

that it was fitted by Mr

it may conveniently be

bisected in C, and at t

CB is attached. D is the

the link AB be guided a

every moment in the ax

For every position C

the angle in a semi-c

scribes a straight line.

This system of links

covery of the Peaucellier-Lipkin linkage.

point can be thrown in

is, parallel to the line j

transferred to the line of

IV. *Hypocycloidal P*

case of the problem of I

cycloidal movement. V

concave circumference

a curve, which is called

diameter of the revolving

VOL. IX.

VOL. IX.




Fig. 6.

MC, with LF, FK form a pantagraph, and for every position of \bar{C} , \bar{P} takes another, equiangular and proportional.

III. *The Gorgon Linkage*.—The parallel motion of Mr Scott Russell is exact, and constitutes a two-bar motion. From the fact that it was fitted by Mr Seaward to the engines of the "Gorgon," it may conveniently be called a Gorgon Linkage. A link AB is bisected in C, and at this point another link CD, equal to CA or CB is attached. D is the centre of revolution. If one end A of the link AB be guided along the axis of x , the motion of B is at every moment in the axis of y from D.

For every position C is the centre of a circle \overline{ADB} , and $\angle ADB$ is the angle in a semi-circle, that is, a right angle. Hence B describes a straight line.

This system of links derives additional interest from the discovery of the Peaucellier cell, as by it the motion of the parallel point can be thrown in a direction at right angles to itself; that is, parallel to the line joining fulcrum and pivot; and this can be transferred to the line of centres by means of the pantagraph.

IV. *Hypocycloidal Parallel Motion*.—Another very interesting case of the problem of Parallel Motion is that produced by a hypocycloidal movement. When one circle is made to revolve on the concave circumference of another circle, any point in it describes a curve, which is called the hypotrochoid or hypocycloid. If the diameter of the revolving circle be equal to the radius of the circle

in which it revolves, the hypotrochoid becomes an ellipse; and if the point be *on* the circumference of the smaller circle, the ellipse degenerates into a straight line.

Let the circle BPC (fig. 7), diameter $BC=AC$, revolve in the circle ABM, radius AC . If the initial position of the describing circle be such that AC




Fig. 7.

be its diameter along the axis of y , the point A will move along the diameter AM . When the circle has moved round to B , let P be the position of A ; then $AB=BP$. Join BKC and KP . Let $ACB = \theta$; $BKP = \phi$. The arc $BP = AB \therefore BK \cdot \phi = AC \cdot \theta = 2 BK \cdot \theta$. Hence $\phi = 2 \theta$. Hence P must lie on AC . (Euc I. 32).

The difficulty of constructing, and the inconvenience experienced in using an arrangement in which the annular wheel comes into action, are so great that it is seldom employed. The following is a simple method of obtaining the hypocycloidal motion without requiring the annular wheel. On the arm ACB (fig. 8),




Fig. 8.

which is made to revolve in a circle MLP round A , attach two toothed wheels LNF and FEM , in gear with each other and with a fixed wheel EOH , whose radius AE is double that of NLF . The centre wheel FEM is used merely for the purpose of reversing the motion of NFL , hence it is immaterial how many teeth it has; the only postulate of the problem

being that LFN revolve on its axis at double the speed of the

arm AC, and in a contrary direction. The number of teeth on HEO is twice the number on NLF. Hence NLF makes two revolutions round its axis, while the arm causes it to revolve round the fixed centre-circle HEO. The axis C is rigidly connected with a wheel XWP, revolving on the upper surface of the plate-arm ABC, whose circumference continually passes through A. Thus while the plate-arm ABC makes one revolution round A in the direction of the hands of a watch, XWPA, which is rigidly connected to the axis C of NLP and partakes of its motion, makes two revolutions in the opposite direction. Thus it revolves round an imaginary annular wheel XYZQ, whose diameter ZQ is double that of its own; and in virtue of hypocycloidal motion, if XWP be the initial position, the point A in the circumference of XWPA will move along ZQ.

In the annexed figure (fig. 9), the *modus operandi* of the machine is sketched. The system is analogous to the sun and planet wheel invented by Watt.




Fig. 9.

It would be going beyond the limits of this paper to direct attention to the two modifications of the Peaucellier cell, which have been called respectively the Quadratic-binomial Extractor, and the Conicograph. The introduction of these into the sphere of mathematical investigation has given several indications of valuable results to be obtained therefrom. For a brief outline of these, we refer the reader to the pamphlet of Professor Sylvester, who not only was the first in this country to direct attention to the general problem, but also had the credit of demonstrating the

higher vantage ground opened to the mathematician by means of Peaucellier's discovery. The extent of our own obligation to him is great.

Not less are we indebted to Professor Kelland, whom we have known both as a teacher and a friend. The valuable hints and suggestions he has given us on this subject we are glad to take this opportunity of acknowledging.

3. Laboratory Notes. By Professor Tait.

- (a.) On the Passage of the Electric Current from Amalgamated Zinc to Zinc Sulphate Solution. By J. G. MacGregor, M.A.
- (b.) On the Thermo-Electric Properties of Cobalt, &c. By Messrs Knott, MacGregor, and C. M. Smith.
- (c.) Measurements of the Potentials required for Long Sparks of a Holtz Machine. By Messrs Macfarlane and Paton.

4. Note on Orthogonal Isothermal Surfaces. By Professor Tait.

5. Notice of some recent Atmospheric Phenomena. By Professor Tait.

6. Report by the Society's Boulder Committee. (Plates II. and III.)


Mr David Milne Home gave in the Third Report of the Society's Boulder Committee, from which the following are extracts:—In November 1875, on the invitation of Sir John Douglas of Glenfinnart, the Convener went to visit him at that place, to have an opportunity of examining several remarkable boulders reported to the Committee as situated in that part of Argyllshire.

1. On the east side of Lochlong, opposite to Ardentinny, there is the farm of Peaton. On this farm, a burn descends from a steepish hill which faces the north. A gneiss boulder lies in a gorge cut by the burn through rocks of clay slate. The boulder




Boulder on Peaton Farm
between rocky sides of a

Loch Long, jammed
Ravine 24 x 16 feet.




Same Boulder viewed from opposite side.



Clach Udelain Boulder of Gneiss 1526 ft.
above Sea, on edge of a cliff about 300 ft high

Fig. 3.




Boulder of Gneiss on a rock of Clay Slate Rock, smoothed from the North, as shown by rough parts facing the South.

Fig. 4.




The Same Boulder, viewed from opposite side. Local name "Giants putting stone."

Fig 5.




Gneiss Boulders North of Knap Farm House in Loch Long resting on smoothed clay slate rocks

Fig 6



Pulag Boulder, near Glenfinnart, on Loch Long, 824 feet above Sea. Its South side rests on a small Boulder

Fig 7.





rests on the rocky sides of the burn, jammed in between the sides. The boulder has some local name like "Jenny Meullen," meaning "House on a knoll." The height of the boulder above the sea is about 226 feet. Its distance from the sea-beach is about three-quarters of a mile. The size of the boulder is about $24 \times 18 \times 12$ feet. It is most probable that the boulder was transported across the loch from the north or north-west, and was arrested in its further progress southward by the hill, on the north side of which it stands.

Two sketches of this boulder are given on Plate II. figs. 1 and 2.

2. Between the site of this boulder and the sea-beach an old sea margin occurs at a height of about 45 feet above the sea (medium level). A number of boulders lie along the line of this sea margin. There is an old sea margin on the opposite or west side of the loch, at exactly the same height—viz., 45 feet.

3. Close to the beach in this part of Lochlong—i.e., about 8 or 9 feet above high-water mark—at a place called "Letter," there lies another gneiss boulder, $12 \times 8 \times 8$ feet. Its long diameter points N.W. by N.—viz., to Glenfinnart Valley.

4. Very near this boulder (about 100 yards to the north) the clay slate rocks have been ground down and smoothed. Their smoothed surfaces show numerous striæ pointing N., 2° or 3° W. (magnetic). The smooth surface dips towards the north at an angle of 3° or 4° .

5. On the hill above Carrick Castle, situated on Lochgoil, there is a boulder called "Clach udelain," or the "Stone nicely balanced." It is at a height of 1526 feet above the sea. This boulder is of gneiss, and lies on rocks of clay slate. It lies on a bare rock, the face of which slopes to N.N.E.—i.e., towards Lochgoil. The boulder is within three or four yards of the edge of a precipitous rocky cliff, which goes vertically down about 500 or 600 feet. The block is of enormous size. Unfortunately the note taken of its dimensions has been lost. This boulder, from its position, could not have fallen from any hill. There is no hill near it from which it could have fallen.

A sketch of this boulder is given on Plate II. fig. 3.

6. The next boulder visited is about two miles to the eastward of the last-mentioned, and is within a quarter of a mile of Lochgoil,

near its junction with Lochlong. It is about 450 feet above the sea. It may be observed, that all the rocks in this district have their smooth faces towards the north, their rough faces towards the south. This boulder has received the name of the "Giant's Putting-Stone," from a legend which alleges that in former times there were giants who inhabited the district on both sides of Lochlong, and who were in the practice of amusing themselves by throwing these huge boulders across the loch. The rock on which it rests slopes gently N. by E. This rock presents a large surface, ground down and smoothed. The space of rock occupied by the superimposed boulder is only 18 inches by 12 inches. It would not be difficult for two men with strong levers to move the boulder from its narrow resting-place, in which case the boulder would probably roll down the steep hillside into the loch.

Two sketches of this boulder are given on Plate III. figs. 4 and 5.

7. To the north of Knap Farm-house, there is a small hill, on or very near the top of which eight or ten boulders are clustered. They suggest the idea that this hill has arrested or interrupted the body, whatever that body was, which transported the boulders, and caused them to be stranded here.

8. There is another hill lower down the valley of Knap (about 480 feet above the sea), the top of which consists of clay slate rocks, rounded and smoothed by some agent passing over from the north. It has received the popular name of the "Pig's Back." Several boulders lie on this ridge. The largest rest on a very small portion of rock.

A sketch of this ridge of rock, with boulders on it, is given on Plate III. fig. 6.

9. Pulag boulder is near the top of a hill to the west of Glenfinnart, about 824 feet above the sea. It is a large block of gneiss, about 7 feet high. There are many other smaller boulders lying near it. The large boulder is almost on the edge of a precipice which goes down at least 200 feet. It could not have been rolled or pushed to its present position. The levels of the district show the greatest openings towards the north—a circumstance which suggests that the boulder came from the north. Moreover, its south end rests on a smaller boulder, which seems to have stopped its progress further south.

A sketch of this boulder is given on Plate III. fig. 7.

10. Along various parts of the hills in this district where their highest ranges are seen against the sky, and at a height of about 2000 feet above the sea, boulders are discernible from a distance, lying on the ridges. It would be very desirable to obtain particular accounts of boulders at so high an elevation.

11. In the last Report of the Committee, notice was taken of a boulder in Ayrshire called the "Hunterston Boulder." Along the same coast, and especially on the property of Mr Alexander of Boydstone, several very large boulders may yet be seen. One, called the "Boydstone Rock or Stone," is situated about two miles north-west of Ardrossan. Some chips of the boulder, sent to the Convener in a letter, show that it is porphyritic. The rocks on this part of the coast are Old Red Sandstone. The boulder is in length about 19 feet, and in breadth about the same. It is partly buried in the mud of the shore. Its highest point is $9\frac{1}{2}$ feet above the shore. It is said to contain 40 cubic yards above the shore line. It is supposed that the boulder is buried to the depth of 5 feet. The tide at high water leaves about 3 feet of the boulder visible. This boulder has inspired the poetic genius of an Ayrshire letter-carrier (Malcolm Kerr, post-messenger between Ardrossan and West Kilbride), who, through Mr Weir of Kirkhall, has sent to the Convener the following stirring address :—

*To the Great Boulder on the shore opposite to the lands of Boydstone,
two miles north of Ardrossan, Ayrshire.*

"Can'st thou speak, old grey stone,
Unto me ?
List thou to the ocean's moan,
I to thee :
Music sweet ! Spirits string
Wild ditties, as they cling
To the big waves which swing
Around thee.

Stranger ! whence didst thou come
To this shore ?
Art thou an Arctic crumb,
Which of yore

On some huge iceberg side
 From thy first home did glide,
 A wanderer on the tide,
 To this shore?

Many eyes with wonder,
 Ages gone,
 Looked on thee! What number
 Yet unknown,
 Will gaze with curious eye,
 Seeking to know thy history,
 And solve a hidden mystery,
 Old grey stone."

Besides the boulder which inspired these verses, there are two others, also on Mr Alexander's property—one of them, as Mr Weir states, "even larger than the big stone at Brigurd, at Hunterston." This larger one it was proposed to split up for building purposes at Ardrossan; but Mr Alexander interfered, and saved it. There are many other boulders along the Ayrshire sea-coast, but none so large as the Boydstone boulder. A chip of one of these sent, shows that it is of gneiss, indicating a northern origin.

12. A report has been received of a gneiss boulder near Dunblane, on the property of Cromlix, belonging to the Hon. Captain Drummond. The length of the boulder is stated to be $17\frac{1}{2}$ feet, its breadth (on an average) 10, its height about 5 feet. Its longer axis lies in a direction south-west and north-east. At its south end it dips into the ground at an angle of 45° . Its weight above ground is estimated at 34 tons. Its height above the sea is about 450 feet. It is about four miles south from the Grampians. The same reporter (Henry Wilkie, Ashfield Works, Dunblane) refers to a group of four boulders in the parish of Redgorton, at the west end of a gravel ridge on the farm of Bertha, the property of Murray Graham, Esq. of Murrayshall. Three of these Redgorton boulders are within an area of 30 yards. They are angular; flat on the top, and some of them square. They seem to be Silurian rocks. They are distant about twelve miles from the Grampians.

13. Notice has been received by the Convener from Mr Robertson, C.E., who is in charge of the Albert New Docks at Leith, that boulder clay was found in excavating for the docks beneath the shore line, to the depth of about 70 feet. The clay was full of

large blocks. Some of these were of sandstone, weighing 10 or 12 tons, and appeared to be of the same rock as that worked at Granton and Craigleith Quarries to the westward. Beneath the boulder clay, there was found what the engineers call a "running sand" lying over strata of shale and sandstone.

Before concluding, the Committee may advert to the circumstance that a part of the district comprehended in this Report was many years ago described by an eminent Scotch geologist—the late Charles M'Laren—and with reference to the very matters embraced in this report. Mr M'Laren read papers in this Society, in the years 1846-47, describing the boulders then existing on the shores of the Gairloch, and on the hills between that loch and Lochlong. Even then the destruction of boulders in that quarter had begun, being appropriated, as Mr M'Laren states, to building purposes; and probably by this time they have all been annihilated. Mr M'Laren in these papers described also the striation and smoothings of the rocks, which he found from the sea-shore up to the tops of the ridges, between the Gairloch and Lochlong, at heights of about 1000 feet above the sea. It is due alike to the memory of our Associate, and to the interests of geological science, to mention, that the boulders referred to by him, as found on the Gairloch, consisted of grey granite, of which he counted above a hundred, one-third of them exceeding 30 tons in weight; as also mica slate, which, though less numerous, had had among them blocks of 60 and 80 tons in weight. As the rocks of the Gairloch *in situ*, are of a more recent kind—viz. clay slate—Mr M'Laren justly inferred that the boulders were of northern origin. For those of granite, he pointed to Ben Cruachan, a mountain exceeding 3000 feet in height, situated to the N.N.W., and distant about thirty miles. The mica slate hills are also in the same direction, somewhat more distant. From his study of the boulders and other phenomena in this district, Mr M'Laren drew two important conclusions. One conclusion was, that the boulders must have been brought to the district from the parent mountains, across valleys and ranges of hills, on ice floating on a sea which stood from 1500 to 2000 feet above the present sea-level, and in which a strong current had prevailed from the N.W. This conclusion, it will be noticed, is confirmed by the facts specified in the

present Report, and also in former Reports by the Committee. The numerous instances given in these Reports of huge boulders shown by their composition to be of northern rocks, clustered frequently on the summits or peaks of hills, at heights of 1500 and 2000 feet above the present sea-level, seem to leave no doubt regarding the soundness of *that* conclusion to which Mr M'Laren had come to thirty years ago.

The other conclusion to which Mr M'Laren came, and which many good geologists of the present day hold, was this, that local glaciers had at one time existed in all those valleys. This he inferred from observing, that the striations on the rocks were all, as he thought, exactly parallel with the axis of the valley, and which here generally runs in a direction north-west and south-east; and also from discovering accumulations of gravel and clay in the form of elongated embankments—some across the valleys, others parallel with the valleys, reminding him of the lateral and terminal moraines of Switzerland.

As to the soundness of this second conclusion, or the correctness of the observations on which it was founded, the Committee give no opinion. None of the members of the Committee have visited the localities, and their function as a Committee has been chiefly to collect facts connected with the boulders. But it may not be out of place to record the fact, that another of our Associates, the late Robert Chambers, who had also given much attention to this branch of geological science, went to examine the districts referred to by Mr M'Laren, and expressed doubts regarding these last-mentioned views. In two papers read by him in this Society in December 1852, Chambers states, that on the rocky ridge between Lochlong and Holy Loch, at a height of 600 feet above the sea, the direction of the striations was not parallel with the valley, but "slanting over the hill;" and as the striations were not merely on the sides of the valleys, but on the tops of the hilly ranges dividing the valleys, he thought they were more probably due to a general glaciation of the country than to local glaciers. Chambers quotes also the opinion of Sir Roderick Murchison, "that there is no imaginable centre for the issue of glaciers of the ordinary kind down the Gairloch." On this point the Committee offer no opinion, and wish only to advert to the investigations of eminent Scotch geolo-

gists who in former days attempted a solution of these difficult questions. The Committee would venture to suggest a re-examination of the localities, as the fuller knowledge now possessed on these subjects may possibly throw a clearer light on the phenomena.

The Committee wish only to add, that if any geologists, whether fellows of this Society or not, happen, in the course of their rambles through the country, to fall in with or hear of any boulders remarkable for size, position, or composition, not yet mentioned in the Committee's Reports, the Convener will be happy to receive a notice of them.

After some conversation, in which Mr Ferguson of Kinmundy, Dr Bryce, the Rev. T. Brown, and others took part, the Report was adopted, and the proceedings terminated.

Donations to the Royal Society Library during Session
1874-75 :—

I. AUTHORS.

- Abbt (Dr Ferdinand). Ueber die Spontanen Rupturen des Herzens. Erlangen, 1875. 8vo.—*From the Author.*
- Armstrong (Sir William) and others. Recent Discussions on the Abolition of Patents for Inventions in the United Kingdom, France, &c. London, 1869. 8vo.—*From R. A. Macfie, Esq.*
- Banner (E. G.). Sanitary Appliances and System of Sanitation. London, 1876. 8vo.—*From the Author.*
- Bartoli (Dottore Adolfo G.). Sopra i Movimenti Prodotti dalla Luce e dal Calore, e sopra il Radiomentio di Crookes. Firenze, 1876. 8vo.—*From the Author.*
- Becker (Franz). Beiträge zur Kenntniss des Tellurs. Erlangen, 1875. 8vo.—*From the Author.*
- Beilby (J. Wood). Reasons suggestive of Mining on Physical Principles for Gold and Coal. Melbourne, 1875. 8vo.—*From the Author.*
- Bidder (Roderich). Ueber Kohleleth's Stellung zum Unsterblichkeitsglauben. Erlangen, 1875. 8vo.—*From the Author.*
- Birnbaum (Rudolph). The Sanitary Powers of the Iron Springs of Schwalbach. Berlin, 1876. 8vo.—*From the Author.*
- Bögler (Dr F.). Klinische Beobachtungen bei Offenbleiben des Foramen Ovale. Erlangen, 1875.—*From the Author.*
- Braun (Wilhelm). Die Singularitäten der Lissajous'schen Stimmgabelcurven. Erlangen, 1875. 8vo.—*From the Author.*
- Bretschneider (Wilhelm). Ueber Curven vierter Ordnung mit drei Doppelpunkten. Stuttgart, 1875. 8vo.—*From the Author.*
- Brimmer (Carl). Ueber die Chemischen Bestandtheile der Angeli-Wurzel. Erlangen, 1875. 8vo.—*From the Author.*
- Bruns (Dr Heinrich). Ueber die Perioden der Elliptischen Integrale. Dorpat, 1875. 4to.—*From the Author.*

- Buchner (Dr Ludwig Andreas). Ueber die Beziehungen der Chemie zur Rechtspflege. Munchen, 1875. 4to.—*From the Author.*
- Burmeister (Dr Hermann). Description physique de la Republique Argentine. Tome Premier. Paris, 1876. 8vo.—*From the Author.*
- Die fossilen Pferde der Pampas-formation. Buenos Aires, 1875. Fol.—*From the Author.*
- Charles (P.). The Geographical Distribution of Animals and Plants in their Wild State. 1876. Salem, Mass. 4to.—*From the Author.*
- Chevreur (M. E.). L'Enseignement devant l'Etude de la Vision. La loi du Contraste Simultané des Couleurs. Paris, 1875. 4to.—*From the Author.*
- Explication de Nombreux Phénomènes qui sont une Conséquence de la Vieillesse. Paris, 1875. 4to.—*From the Author.*
- Etudes des Procédés de l'Esprit Humain dans la Recherche de l'Inconnu à l'aide de Observation et de l'Experience, et du moyen de savoir s'il a trouvé l'Erreur ou la Verité. Paris, 1874. 4to.—*From the Author.*
- Croizier (Le C. De). La Perse et les Persans. Paris, 1873. 8vo.—*From the Author.*
- Day (St John V.). A Paper on Recent Arrangements of Continuous Brakes. 1875. 8vo.—*From the Author.*
- Diehl (Dr G.). Beiträge zur Pathologie und Therapie des Diabetes Mellitus. Erlangen, 1875. 8vo.—*From the Author.*
- Disse (Dr Joseph). Beiträge zur Anatomie des Menschlichen Kehlkopfs. Bonn, 1875. 8vo.—*From the Author.*
- Dana (Prof. A.). Il Passaggio di Venere sul Sole, Osservato a Muddapur. Palermo, 1875. 4to.—*From the Author.*
- L'Aneroide a Vite Micrometrica Sperimentata colle differenze di livello della Strada Ferrata delle Alpi. Torino, 1875. 8vo.—*From the Author.*
- Eidam (Dr H.). Ueber die Bildungsweise intraperitonealer tumoren im Kleinen Becken. Leipzig, 1875. 8vo.—*From the Author.*

- Ennis (John), A.M. The Origin of the Stars, and the Causes of their Motions and their Light. London, 1876. 8vo.—*From the Author.*
- Faber (Dr J.). Die Embolie der Arteria Mesenterica superior. Erlangen, 1875. 8vo.—*From the Author.*
- Fleischmann (Johann Karl). Kritische Studien über die Kunst der Charakteristik bei Æschylos und Sophokles. Nurnberg, 1875. 8vo.—*From the Author.*
- Forster (Dr S. von). Ueber congenitale Hypoplasie des Uterus. Erlangen, 1875. 8vo.—*From the Author.*
- Friertag (Isaac). Ueber die Bildung der Hare. Dorpat, 1875. 8vo.—*From the Author.*
- Fries, E. Icones Hymenomycetum. Fasc. 7-10. 4to.—*From the Royal Academy of Sciences, Stockholm.*
- Furbringer (Dr Max). Beitrag zur Kenntniss der Kehlkopfmuskulatur. Jena, 1875. 8vo.—*From the Author.*
- Garstin (J. R.). Facts and Reasonings adduced in support of the course taken by the Royal Irish Academy respecting the recent action of certain Government Departments. Dublin, 1876. 8vo.—*From the Author.*
- Geer (Louis de). Minnestekning öfver Hans Järta. Stockholm, 1874. 8vo.—*From the Author.*
- Gerichten (Dr E. Von). Die Theorie der Säuren und Salzbildung und die Electrochemische Theorie. Erlangen, 1875. 8vo.—*From the Author.*
- Thesen welche zur Erlangung der Venia Legendi in der Philosophischen Facultat, &c. Erlangen. 1875. 4to.—*From the Author.*
- Gordan (Dr Paul). Ueber das Formensystem binärer Formen. Leipzig, 1875. 8vo.—*From the Author.*
- Gresbeck (Berthold). Ursache der Irritation cerebri infantum und deren Behandlung. Erlangen, 1875. 8vo.—*From the Author.*
- Grünewald (Max). Ueber den Kaiserschnitt. Munchen, 1876. 8vo.—*From the Author.*
- Hamilton (H.). Minnesteckning öfver Jacob August von Hartmansdorff. Stockholm, 1872. 8vo.—*From the Author.*

- Harneck (Axel.). Ueber die Verwerthung der elliptischen functionen für die Geometrie der Curven dritten Grades. Leipzig, 1875. 8vo.—*From the Author.*
- Hartley (Walter Noel). Air, and its Relations to Life. London, 1875. Second edition, 1876. 8vo.—*From the Author.*
- Hartung (Dr E.). Ueber einen Fall von Mamma accessoria. Erlangen, 187. 8vo.—*From the Author.*
- Hayter (H. H.). Victorian Year-book for 1874. London, 1875. 8vo.—*From the Author.*
- Herzog (Wilhelm). Beiträge zur Näheren Kenntniss der Structur der Seline. Erlangen, 1875. 8vo.—*From the Author.*
- Hess (Dr Otto). Ueber einen Fall von Heterotaxie der Brust- und Baueingeweide. Erlangen, 1875. 8vo.—*From the Author.*
- Hoerner (Dr C.). De extremo Græcorum discrimine quomodo in Iliade descriptum sit. Erlangen, 1875. 8vo.—*From the Author.*
- Hooker (Dr J. D.), C.B. The Flora of British India. Part IV. 8vo.—*From the India Office.*
- Hornberger (Richard). Einige Verbindungen des Zirkoniums. Erlangen, 1875. 8vo.—*From the Author.*
- Hornstein (Carl). Astronomische Magnetische und Meteorologische beobachtungen an der K. K. Sternwarte zu Prag im Jahre, 1875-76. 4to.—*From the Author.*
- Jakowicki (Anton). Zur Physiologischen Wirkung der Bluttransfusionen. Dorpat, 1875. 8vo.—*From the Author.*
- Johanson (Edwin). Beiträge zur Chemie der Eichen-Weiden und Ulmennride. Dorpat, 1870. 8vo.—*From the Author.*
- Kaulbars (Franz). Trophische und vasmotorische Störungen peripherer Nervenverletzungen. Jena, 1874. 8vo.—*From the Author.*
- Kiesselbach (Wilhelm). Beiträge zur Näheren Kenntniss der Sogenannten Grauen Degeneration des Sehnerven bei Erkrankungen des Cerebrospinalsystems. Erlangen, 1875. 8vo.—*From the Author.*
- Kuhn (Moriz). Ueber die Beziehung Zwischen Druck, Volumen und Temperatur bei Gasen. Wein, 1875. 4to.—*From the Author.*

- Lassel (W.). On Publishing the Specula of Reflecting Telescopes. 1874. 4to.—*From the Author.*
- Leuckart (Rudolf). Die Menschlichen Parasiten und die von Ihnen Herrührenden Krankheiten. 1876. 8vo.—*From the Author.*
- Luvini (Jean). Le Diéthéroscope. Turin, 1876. 8vo.—*From the Author.*
- Lyman (Theodore). Commemorative Notice of Louis Agassiz. 8vo.—*From the Author.*
- Macdonald (James). The Origin and Destiny of Comets. Prah-
ran, 1869. 8vo.—*From the Author.*
- Maison (Dr Otto). Ueber die Resection des Unterkiefers. Er-
langen, 1875. 8vo.—*From the Author.*
- Mandelstam (Dr N.). Ueber den Einfluss Chemischer Agentien
auf die Erregbarkeit der Nerven. Erlangen, 1875. 8vo.—
From the Author.
- Mazzola (Giuseppe). Effemeridi del Sole, della Luna. Torino,
1874-75. 8vo.—*From the Author.*
- Mehlis (Christian). Die Grundidee des Hermes von Standpunkte
der Vergleichenden Mythologie. Erlangen, 1875. 8vo.—
From the Author.
- Meissner (Dr A.). Ein Fall von Verengerung der Pfortader durch
chronische Peritonitis bei Gallensteinerkrankung. Erlangen,
1875. 8vo.—*From the Author.*
- Michel (Dr J.). Die Histologische Structur des Irisstroma. Er-
langen, 1875. 8vo.—*From the Author.*
- Molitor (Eduard). Die Behandlung des Fiebers bei Typhus
und Pneumonie. Munchen, 1875. 8vo.—*From the
Author.*
- Monteiro (Joachim John). Angola and the River Congo. Vols.
I. and II. London, 1875. 8vo.—*From the Author.*
- Müller (Baron Fredinandus von). Fragmenta Phytographiæ.
Vol. IX. Melbourne. 8vo.—*From the Author.*
- Descriptive Notes on Papuan Plants. Melbourne. 8vo.—
From the Author.
- Naumann (Alexander). Jahresbericht über die Fortschritte der
Chemie für 1873. Heft 3. 8vo.—*From the Editor.*

- Nussbaum (Heinrich). Beiträge zur Kenntniss der Anatomie und Physiologie der Herznerven und zur Physiologischen Wirkung des Curare. Dorpat, 1875. 8vo.—*From the Author.*
- Oudemans (Dr J. A. C.). Die Triangulation von Java Ausgeführt vom Personal des Geographischen Dienstes in Niederländisch Ost-Indien. Batavia, 1875. 4to.—*From the Author.*
- Paolo (Riccardi). Annuario della Societa dei Naturalisti in Modena, 1875. Anno IX., X. 8vo.—*From the Author.*
- Paulus (Franz). Verhartung und Verknocherung des Haematoms der Dura mater und ihre Beziehung zur Heilung desselben. Erlangen, 1875. *From the Author.*
- Penzoldt (Dr F.). Die Magenerweiterung. Erlangen, 1875. 8vo.—*From the Author.*
- Pfeuffer (Dr P.). Experimentelle beiträge zur Resorption gefärbter fette.—Erlangen, 1875. 8vo.—*From the Author.*
- Pleyte (W.). Les Peintures murales de l'Eglise dédiée a Saint Jaques à Utrecht. 1874. Fol.—*From the Author.*
- Popp (Friedrich). Ueber Embolie der Arteria centralis retinae. Regensburg, 1875. 8vo.—*From the Author.*
- Quetelet (M. Ern). La Tempete du 12 Mars. 1876. 8vo.—*From the Author.*
- Raab (Dr Jos.). Die Tetanie. Erlangen, 1875. 8vo.—*From the Author.*
- Ribot (F.). Revue Philosophique de la France et de l'Etranger. Paris, 1876. 8vo.—*From the Author.*
- Richardson (Ralph). The Ice Age in Britain. Edinburgh, 1876. 8vo.—*From the Author.*
- Rothhaupt (F. R.). Die Pulsformen der Paralysis progrediens. Mellrichstadt, 1874. 8vo.—*From the Author.*
- Sandon (Lord). Letter to the President of the Royal Irish Academy. 1876. 8vo.—*From the Royal Irish Academy.*
- Schmidt (Alfred). Das Russische Geldwesen Während der Finanzverwaltung des Grafen Cancren von 1823-1844. St Petersburg, 1875. 8vo.—*From the Author.*
- "Searcher," a Truth in Extremis; A Plea against the Murder of Science by the Gold Poison. Oxford, 1876. 8vo.—*From the Author.*

Steinmann (F.). Ueber Retroversionen und Retroflektionen der Gebärmutter und deren Behandlung. Erlangen, 1875. 8vo.
—*From the Author.*

Steubing (Dr W.). Ueber des Schrägverengte Becken und seinen Einfluss auf die Geburt. Erlangen, 1875. 8vo.—*From the Author.*

Stewart (C. M. A.). International Correspondence by means of Numbers. London, 1874. 8vo.—*From the Author.*

Sudhoff (Karl). Ueber das Primäre Multiple Carcinome des Knochensystems. Erlangen, 1875. 8vo.—*From the Author.*

Traub (Dr M.). Driemaandelijksch Botanisch Literatuurverzicht. Nijmegen, 1874. 8vo.—*From the Author.*

Trumpp (Ernst). Einleitung in das Studium der Arabischen Grammatiker. München, 1876. 8vo.—*From the Author.*

Veh (Friedrich). Ueber die Wirksamkeit klar filtrirter faulender Flüssigkeiten. Dorpat, 1875. 8vo.—*From the Author.*

Vincent (Charles W., F.R.S.E.). The Year-Book of Facts in Science and the Arts for 1875. London. 8vo.—*From the Author.*

Vinchon-Thiesset (Artus). La Cause des Effets. Saint Quentin, 75. 8vo.—*From the Author.*

Vogt (Dr Günther). Zur Statistik und Casuistik der Progressiven Muskelatrophie. Erlangen, 1875. 8vo.—*From the Author.*

Wedekind (L.). Beiträge zur Geometrischen interpretation Binärer Formen. Erlangen, 1875. 8vo.—*From the Author.*

Wikszemski (Adam). Beiträge zur Kenntniss der giftigen Wirkung des Wasserschiefelings. 8vo.—*From the Author.*

Zielonko (Justus). Pathologisch-Anatomische und experimentelle Studien über Hypertrophie des Herzens. Dorpat, 1875. 8vo.
—*From the Author.*

II. TRANSACTIONS AND PROCEEDINGS OF LEARNED SOCIETIES, ACADEMIES, ETC.

Amsterdam.—Flora Batava. Nos. 227-231. 4to.—*From the King of Holland.*

Jaarboek van de Koninklijke Akademie van Wetenschappen gevestigd te Amsterdam voor 1774. 8vo.—*From the Academy.*

Amsterdam.—Processen-verbaal van de Gewone Vergaderingen der Koninklijke Academie van Wetenschappen, 1874-1875. 8vo.—*From the Academy.*

Verhandelingen der Koninklijke Academie van Wetenschappen. Deel VIII.—XV. 4to.—*From the Academy.*

Verslagen en Mededeelingen der Koninklijke Academie van Wetenschappen. Afdeling Naturkunde. Deel IX. 8vo.—*From the Academy.*

Baltimore.—First Annual Report of the Provost to the Trustees of the Peabody Institute. 1868.

Ninth Annual Report (1876) of the Peabody Institute. 8vo.—*From the Institute.*

Basel.—Verhandlungen der Naturforschenden Gesellschaft. Sechster Theil, Zweites Heft. 8vo.—*From the Society.*

Berlin.—Abhandlungen der Königl. Akademie der Wissenschaften zu Berlin. 1874-1875. 4to.—*From the Academy.*

Die Fortschritte der Physik im Jahre 1870. Dargestellt von der Physikalischen Gesellschaft zu Berlin. XXVI. Jahrgang, Abth. 1-2; 1871, XXVI. Jahrgang, Abth. 1-2. 8vo.—*From the Society.*

Jahresbericht der Commission zur Wissenschaftlichen Untersuchung der Deutschen Meere in Kiel für die Jahre 1872-73, 2 und 3 Jahrgang. Fol.—*From the Commission.*

Monatsbericht der Königl. Preussischen Akademie der Wissenschaften zu Berlin. 1875—Juni, Juli, August, September, October, November, December. 1876—Januar, Februar, März, April, Mai. 8vo.—*From the Academy.*

Berne.—Mittheilungen der Naturforschenden Gesellschaft in Bern aus dem Jahre 1874. Nos. 828-873, 878-905. 8vo.—*From the Society.*

Bombay.—Census of the Bombay Presidency, taken on the 21st February 1872. Part 3d. 4to.—*From the Bombay Government.*

The Indian Antiquary. Vol. IV. from September to December 1875; Vol. V. from January to July 1876. 4to.—*From the Publishers.*

- Bombay*.—Journal of the Bombay Branch of the Royal Asiatic Society, 1875. Vol. XI. Nos. 31, 32. 8vo.—*From the Society*.
- Bonn*.—Verhandlungen des Naturhistorischen Vereines der Preussischen Rheinlande und Westfalens. Jahrgang XXXI. Folge 3; XXXII. Folge 4. 8vo.—*From the Society*.
- Bordeaux*.—Mémoires de la Société des Sciences Physiques et Naturelles de Bordeaux. Tome I. Second Series; Second Caher. 8vo.—*From the Society*.
- Boston*.—Annual Report of the Trustees of the Museum of Comparative Zoology at Harvard College in Cambridge 1874–1875. 8vo.—*From the Trustees*.
- Appalachia*. Vol. I. No. 1. 8vo.—*From the Appalachian Mountain Club*.
- Memoirs of the Boston Society of Natural History. Vol. II. Part 4; Nos. 2–4. 4to.—*From the Society*.
- Proceedings of the Society of Natural History. Vol. XVII. Parts 3, 4; XVIII. Parts 1, 2. 8vo.—*From the Society*.
- Occasional Papers of the Boston Society of Natural History. 8vo.—*From the Society*.
- Proceedings of the American Academy of Arts and Sciences. New Series, Vol. II. 8vo.—*From the Academy*.
- The Complete Works of Count Rumford. Published by the American Academy of Arts and Sciences. Vol. IV.—*From the Academy*.
- Brussels*.—Bulletin de l'Académie Royale des Sciences, des Lettres et des Beaux-Arts de Belgique. Tome XII. Partie 1; XXXI. No. 5; XXXII. No. 11; XL. Nos. 7–12; XLI. Nos. 1–6. 8vo.—*From the Academy*.
- Mémoires de l'Académie Royale des Sciences, des Lettres et des Beaux-Arts de Belgique. Tome XLI. Part 1, 2. 4to.—*From the Academy*.
- Mémoires Couronnés et Mémoires des Savants Etrangers Publiées par l'Académie Royale des Sciences, des Lettres et des Beaux-Arts de Belgique. Tome XXXVIII., XXXIX. Part 1. 4to.—*From the Academy*.
- Mémoires Couronnés et Autres, Mémoires Publiées par l'Académie Royale des Sciences, des Lettres et des Beaux-Arts de Belgique. Tome XXIV., XXV., XXVI. 8vo.—*From the Academy*.

Brussels.—Biographie Nationale publié par l'Académie des Sciences, des Lettres et des Beaux-Arts de Belgique. Tome Cinquieme, Part 1. 8vo.—*From the Academy.*

Notices Biographiques et Bibliographiques. 1874. 8vo.—*From the Academy.*

Annuaire de l'Académie Royale des Sciences, des Lettres et des Beaux-Arts de Belgique. 1876. 8vo.—*From the Academy.*

Budapest.—A Magyar Tudományos Akadémia Ertesitoje A. M. T. Akadémia Rendelelebol Szerkeszti A Fotitkar. 1873, Szam 8-11; 1874, Szam 1-14-17. 8vo.—*From the Academy.*

Mathematekaies Termeszett-udományi Közlemenyek vonatkozólag Aszlavia Viszonyokra Kiadja A Magyar Tudományos Akadémia. Kotel VII., VIII., IX., X. 8vo.—*From the Academy.*

Icones Selectæ Hymenomycetum Hungariæ Magyar-ország Harty Agombainak Valogatott Kepel. No. 2, 3, 7-10. Fol.—*From the Academy.*

Az Emberi vese Visszere-Rendszere Dr Senhossek Jozsef. Evk. XIV. Kot. 4. 4to.—*From the Academy.*

Adatok à Szem Perczhartyaja Szövet es Elettanahoz Dr Hanhoffer Lajos, Evk. XIV. Kot. 5. 4to.—*From the Academy.*

Ertezesek A Mathemat, Tudományok, Korebol Kiadja A Magyar Tudományos Akadémia. A III. Osztaly Rendelelebol Szerkeszti Szabo Jozsef Osztalytkar. 1873, Kotel II. Szam 3-6; 1874, III. Szam 1-8, 15; 1873, IV.; 1873, Szam 3-6; IV. 1875, Szam 1-3; V. 1874, Szam 1-9; VI. 1875, Szam 1-6. 8vo.—*From the Academy.*

M. Tudom Akadémiai Almanach Ossillagaszati es Kozen-seges. 1873-75. 8vo.—*From the Academy.*

Buenos Aires.—Acta de la Academia Nacional de Ciencias exactas existente en la Univsraidad de Cordova. Tom I. 4to.—*From the Author.*

Calcutta.—Memoirs of the Geological Survey of India. Vol. XI. part 2, 8vo. Palæontologia, Vol. I. Parts 2-4. 4to.—*From the Survey.*

- Calcutta*.—Records of the Geological Survey of India. Vol. VIII. Parts 1-4. 8vo.—*From the Survey.*
- Journal of the Asiatic Society of Bengal. 1875, Nos. 2-4, Part 1; Part 2, No. 1-3, extra No. 8vo.—*From the Society.*
- Proceedings of the Asiatic Society of Bengal. 1875, Nos. 6-10; 1876. Nos. 1, 2. 8vo.—*From the Society.*
- Report of the Meteorological Reporter to the Government of Bengal for the years 1867 to 1874. By W. G. Wilson, M.A.L.C.E. Fol.—*From the Government of Bengal.*
- Report of the Midnapore and Burdwan Cyclone of the 15th and 16th of October 1874. By W. G. Wilson, M.A.L.C.E. Fol.—*From the Government of Bengal.*
- Report of the Commissioners appointed to Inquire into the Origin, Nature, &c., of the Indian Cattle Plagues, with Appendices. 1871. Fol.—*From the Commissioner.*
- California*.—Proceedings of the California Academy of Sciences. Vol. V. Part 3. 8vo.—*From the Academy.*
- Cambridge (U.S.)*.—Catalogus Universitatis Harvardianæ. 1875. 8vo.—*From the University.*
- The Harvard University Catalogue. 1874-75. 8vo.—*From the University.*
- Forty-Ninth Annual Report of the President of Harvard College. 1873-74. 8vo.—*From the College.*
- The Treasurer's Statement of Harvard College. 1874. 8vo.—*From the College.*
- Illustrated Catalogue of the Museum of Comparative Zoology at Harvard College. No. VIII. Part 2. 4to.—*From the College.*
- Canada*.—Report of the Progress of the Geological Survey of Canada for 1874-75. 8vo.—*From the Geological Survey.*
- Cape of Good Hope*.—Observations made at the Magnetical and Meteorological Observatory at the Cape of Good Hope. Vol. II. 1841-1846. 4to.—*From Her Majesty's Government.*
- Catania*.—Atti dell Accademia Gioenia dé Scienze Naturali de Catania. Tomo VI., IX. 4to.—*From the Academy.*

- Cherbourg*.—Mémoires de de la Société Nationale des Sciences Naturelles de Cherbourg. Tome IX. 8vo.—*From the Society.*
- Chur*.—Verhandlungen der Schweézerischen Naturforschenden Gesellschaft in Chur. Jahresbericht 1873-74. 8vo.—*From the Society.*
- Copenhugen*.—Oversigt over det Kongelige danske Videnskabernes Selskabs Forhandlingar og dets Medlemmers Arbejder i, Aaret 1874, No. 3; 1875, No. 1. 8vo.—*From the Society.*
- Mémoires de l'Académie Royale de Copenhagen. 5th Serie, Vol. X. No. 7-9; Vol. XI. No. 1, 2; XII. No. 1, 2. 4to.—*From the Academy.*
- Dehra Doon*.—General Report on the Operations of the Great Trigonometrical Survey of India during 1874-75. Fol.—*From the Survey.*
- Dublin*.—Journal of the Geological Society. Vol. I. Part 1; II. Parts 3, 4. 8vo.—*From the Society.*
- Journal of the Royal Dublin Society. Vol. VII. Part 1. 8vo.—*From the Society.*
- Proceedings of the Royal Irish Academy. Vol. II. Nos. 1-3. 8vo.—*From the Academy.*
- Transactions of the Royal Irish Academy (Science). Vol. XXV. Parts 10-18. 4to.—*From the Academy.*
- Edinburgh*.—Proceedings of the Royal Physical Society. Session 1874-75. 8vo.—*From the Society.*
- Transactions of the Highland and Agricultural Society of Scotland. Vol. VIII. 8vo.—*From the Society.*
- Catalogue of the Printed Books in the Library of the Faculty of Advocates. Vol. IV. 4to.—*From the Library.*
- Transactions of the Royal Society of Arts. Vol. IX. Parts 2, 3. 8vo.—*From the Society.*
- Transactions and Proceedings of the Botanical Society. Vol. XII. Part 2. 8vo.—*From the Society.*
- Forty-Eighth Annual Report of the Council of the Royal Scottish Academy of Painting, Sculpture, and Architecture. 1875. 8vo.—*From the Academy.*

Edinburgh.—Journal of the Scottish Meteorological Society. Nos. 47–48. 8vo.—*From the Society.*

Erlangen.—Sitzungsberichte der Physicalisch-Medizinischen Societät zu Erlangen. Heft 6. 8vo.—*From the Society.*

Frankfort.—Abhandlungen herausgegeben von der Senckenbergischen Naturforschenden Gesellschaft. Band IX. Heft 3, 4; X. Heft 1–4. 4to.—*From the Society.*

Bericht über die Senckenbergische Naturforschende Gesellschaft. 1873–74; 1874–75. 8vo.—*From the Society.*

Geneva.—Mémoires de la Société de Physique et d'Histoire Naturelle de Genève. Tome XXIV. Part 1. 4to.—*From the Society.*

Glasgow.—Proceedings of the Philosophical Society. 1875–76. 8vo.—*From the Society.*

Gottingen.—Abhandlungen der Königlichen Gesellschaft der Wissenschaften zu Gottingen. Band XX. 4to.—*From the Society.*

Nachrichten von der K. Gesellschaft der Wissenschaften und der Georg-Augusts-Universität, aus dem Jahre 1875. 12mo.—*From the University.*

Gratz.—Mittheilungen des Naturwissenschaftlichen Vereines für Steiermark. Jahrgang 1875. 8vo.—*From the Society.*

Greenwich.—Astronomical and Magnetical Observations made at the Royal Observatory in the year 1873. 4to.—*From the Observatory.*

Haarlem.—Archives Néerlandaises des Sciences Exactes et Naturelles publiées par la Société Hollandaise à Haarlem. Tome X. Liv. 1–5; XI. Liv. 1–3. 8vo.—*From the Society.*

Archives du Musée Teyler. Vol. IV. Fasc. 1. 8vo.—*From the Society.*

Naturerkundige Verhandelingen der Hollandsche Maatschappij der Wetenschappen. Deel II. No. 5. 4to.—*From the Society.*

Hamburg.—Verhandlungen des Vereines für Naturwissenschaftliche Unterhaltung zu Hamburg. 1875. 8vo.—*From the Society.*

Helsingforsia.—Acta Societatis Scientiarum Fennicæ. Tome X. 1875. 4to.—*From the Society*.

Innsbruck.—Berichte des Naturwissenschaftlich - Medizinischen Vereines in Innsbruck. Jahrgang VI. 1875, Heft 1. 8vo.—*From the Society*.

Jena.—Jenaische Zeitschrift für Naturwissenschaft herausgegeben von der Medicinisch-Naturwissenschaftlichen Gesellschaft zu Jena. Band IX. Heft 4; X. Heft 1-3. 8vo.—*From the Society*.

Kasan.—Reports of the University of Kasan. 1875. Nos. 1-6. 8vo.—*From the University*.

Kiel.—Schriften der Universität zu Kiel aus dem Jahre 1874. Band XXI., XXII. 4to.—*From the University*.

Leeds.—Annual Report of the Leeds Philosophical and Literary Society. 1874-75. 8vo.—*From the Society*.

Leiden.—Annalen der Sternwarte, herausgegeben von Dr H. G. Van de Sande Bakhuyzen Verter. 4to.—*From the Observatory*.

Leipzig.—Ueber die Darstellung der Geraden Aufsteigung des Mondes in Function der Länge in der Bahn und der Knotenlänge. Band X. No. 8. P. A. Hansen. 8vo.—*From the Royal Saxon Academy*.

Dioptrische Untersuchungen mit Berücksichtigung der Farbenzerstreuung und der Abweichung wegen Kugelform. P. A. Hansen. Band X. No. 9. 8vo.—*From the Royal Saxon Academy*.

Ueber das Aelius und Sabinus System wie über Einige Verwandte Rechts-Systeme von Moritz Voigt. Band VII. No. 4. 8vo.—*From the Royal Saxon Academy*.

Von der Bestimmung der Theilungsfehler eines Gradlinigen Maassstabes. P. A. Hansen. Band X. No. 7. 8vo.—*From the Royal Saxon Academy*.

Preisschriften gekrönt und herausgegeben von der Fürstlich Jablonowskischen Gesellschaft zu Leipzig. 8vo.—*From the Royal Saxon Academy*.

Ueber die Störungen der Grossen Planeten insbesondere des Jupiter. P. A. Hansen. Band XI. No. 4. 8vo.—*From the Royal Saxon Academy*.

Leipzig.—Elektrische Untersuchungen über die Thermaelektrischen Eigenschaften Des Gypses des Diopsids des Orthoklases, des Albits und des Periklins. W. G. Hanket. Band XI. Nos. 3, 5. 8vo.—*From the Royal Saxon Academy.*

Bericht über die Verhandlungen der Königlich. Sächsischen Gesellschaft der Wissenschaften zu Leipzig. Math. Phys. Classe, 1873, Nos. 3-7; 1874, Nos. 1-5; 1875, Nos. 1. Phil.-Hist. Classe, 1873; 1872, Nos. 1, 2; 1871, No. 1. 8vo.—*From the Royal Saxon Academy.*

Ueber den Ausgangswerth ker Kleinsten Abweichungssumme Dressen Bestimmung Verwendung und Verallgemeinerung. G. Th. Fechner. Band XI. No. 1. 8vo.—*From the Royal Saxon Academy.*

Ueber das von Weber für die Elektrischen Kräfte Aufgestellte Gesetz. Carl Neumann. Band XI. No. 2. 8vo.—*From the Royal Saxon Academy.*

Erster Jahresbericht der Zoologischen Station in Neapel. 8vo.—*From the Society.*

Die Geschichtschreibung über den Schmalkaldischen Krieg, von Georg Voigt. Band VI. No. 6. 8vo.—*From the Royal Saxon Academy.*

Die Epheten und der Areopag vor Solon von Ludvig Lange. Band VII. No. 2. 8vo.—*From the Royal Saxon Academy.*

Zur Charakteristik König Johann's von Sachsen in Seinen Verhältniss zu Wissenschaft und Kunst, von Dr Johann Paul, von Falkenstein. Band VII. No. 3. 8vo.—*From the Royal Saxon Academy.*

Liverpool.—Proceedings of the Literary and Philosophical Society, No. 29. 8vo.—*From the Society.*

London.—Calendar of the Royal College of Surgeons of England. 1875. 8vo.—*From the College.*

Report of the Meteorological Committee to the President and Council of the Royal Society. 1875. 8vo.—*From the Meteorological Committee of the Royal Society.*

Contribution to the Meteorology of Japan, by Staff-Commander Thomas H. Lizard, H.M.S. "Challenger." 4to.—*From the Meteorological Committee of the Royal Society.*

London.—On the Physical Geography of the part of the Atlantic which lies between 20° N. and 10° S., and extends from 10° to 40° W. By Captain Toynbee, F.R.A.S., F.R.G.S.

8vo.—*From Meteorological Committee of the Royal Society.*

Statistical Report on the Health of the Navy for the year 1874. 8vo.—*From the Admiralty.*

H.M.S. "Challenger" Report on Ocean Soundings and Temperatures—Pacific Ocean, No. 5; H.M.S. "Valorous" Deep-Sea Soundings and Temperatures—North Atlantic Ocean. 1875. Fol.—*From the Lords Commissioners of the Admiralty.*

Memoirs of the Royal Astronomical Society. Vol. XLII. 1873-75. 4to.—*From the Society.*

Proceedings of the Royal Society. Vol. XXIII. Vol. XXIV. Nos. 164-170; XXV. No. 171. 8vo.—*From the Society.*

Transactions of the Royal Society. Vol. CLXV. Part 1. List of Fellows, 1874, 4to.—*From the Society.*

Quarterly Weather Report of the Meteorological Office. 1874, Part 2. 8vo.—*From the Meteorological Committee of the Royal Society.*

Results of the Monthly Observations of Magnetic Dip, Horizontal Force, and Declination, made at the Kew Observatory, from April 1869 to March 1875. By the Kew Committee. 8vo.—*From the Royal Society.*

Transactions of the Clinical Society. Vol. VIII. 8vo.—*From the Society.*

Transactions of the Pathological Society. Vol. XXVI. 8vo.—*From the Society.*

Transactions of the Linnean Society. Second Series (Botany), Vol. I. Part 2; Second Series (Zoology), Vol. I. Part 2. 4to.—*From the Society.*

The Anatomy of the Lymphatic System. By E. Klein, M.D.—The Lung, II. 8vo.—*From the Royal Society.*

Proceedings of the Chemical Society. Vol. XIII., October, November, December; XIV., January, February, March, April, May, June, July. 8vo.—*From the Society.*

Proceedings of the Royal Geographical Society. Vol. X. Nos. 1, 2; Vol. XX. Nos. 3-6. 8vo.—*From the Society.*

- London.*—Journal of the Statistical Society. Vol. XXXVIII. Part 4; Vol. XXXIX. Parts 1, 2. 8vo.—*From the Society.*
- Journal of the Linnean Society. Botany, Vol. XV. Nos. 81–84; Zoology, Vol. XII. No. 60–63. List of Members. 8vo.—*From the Society.*
- Journal of the Geological Society. Vol. XXXI. Part 4; XXXII. Parts 1, 2. List of Members. 8vo.—*From the Society.*
- Transactions of the Linnean Society. Second Series. (Botany), Vol. I. Part 3; (Zoology), Vol. I. Part 3. 8vo.—*From the Society.*
- Proceedings of the Linnean Society of the Session 1874–75. 8vo.—*From the Society.*
- Journal of the Royal Asiatic Society of Great Britain and Ireland. Vol. VIII. Part 1. 8vo.—*From the Society.*
- Proceedings of the Royal Institution of Great Britain. Vol. IV. Parts 7, 8; Vol. VII. Parts 5, 6. List of Members. 8vo.—*From the Society.*
- Proceedings of the Geological Association. Vol. IV. Nos. 4–7. Annual Reports, 1875. 8vo.—*From the Association.*
- Proceedings of the Mathematical Society. Nos. 83–92. 8vo.—*From the Society.*
- Proceedings of the Society of Antiquaries. Vol. VI. No. 5. 8vo.—*From the Society.*
- Journal of the Royal Institution. Vol. V. No. 27; VI. No. 28.—8vo.—*From the Institution.*
- Transactions of the Zoological Society. Vol. IX. Parts 5–7. 4to.—*From the Society.*
- Proceedings of the Zoological Society, 1875. Parts 2, 3. 8vo.—*From the Society.*
- Revised List of the Vertebrated Animals now or lately living in the Gardens of the Zoological Society of London. Supplement. 8vo.—*From the Zoological Society.*
- Transactions of the Royal Medical and Chirurgical Society. Vol. LVIII. 8vo.—*From the Society.*
- Proceedings of the Royal Medical and Chirurgical Society. Vol. VIII. Nos. 1, 2. 8vo.—*From the Society.*

- London*.—Journal of the East India Association. Vol. IX
Nos. 4, 5. 8vo.—*From the Association*.
Journal of the Royal Geographical Society. Vol. XLV.
8vo.—*From the Society*.
Proceedings of the Institution of Civil Engineers. Vols.
XLII., XLIII., Parts 1, 2. 8vo.—*From the Institution*.
Lucern.—Verhandlungen der Schweizerischen Naturforschenden
Gesellschaft in Andermatt den 12, 13, und 14 September
1875. 8vo.—*From the Society*.
Lund.—Acta Universitatis Lundensis Lunds Universitets Ars-
Skrift Mathematik och Naturvetenskap för år 1871-73.
Tom. IX., X.—Philosophi Spakvetenskap och Historia.
1872-73—Tom. IX., X. 1871—Theologie. 4to.—*From*
the University.
Madrid.—Memorias de la Real Academia de Ciencias Exactas,
Físicas y Naturales de Madrid. Tomo VI. Parts 1,
4to.—*From the Academy*.
Manchester.—Transactions of the Geological Society. Vol. XIV
Parts 1-3. 8vo.—*From the Society*.
Proceedings of the Literary and Philosophical Society. Vol.
XIV. Nos. 11, 12. 8vo.—*From the Society*.
Mexico.—Boletén de la Sociedad de Geografía-y-Estadística de la
República Mexicana. Tomo II. Nos. 5-7. 8vo.—*From*
the Society.
Montpellier.—Mémoires Académie des Sciences et Lettres de Mont-
pellier; Section des Sciences. Tome VI. Fasc. 2, 3; Tome
VII., VIII. Fasc. 1, 2. De la Section de Médecine—Tome
IV. Fasc. 3-6. De la Section des Lettres—Tome IV.
Fasc. 2-4. Tome V. Plates. 4to.—*From the Academy*.
Moscow.—Bulletin de la Société Impériale des Naturalistes 1875,
Nos. 1-4; 1876, No. 1. 8vo.—*From the Society*.
Munich.—Catalogus Codicum Latinorum Bibliothecæ Regiæ
Monacensis. Tomi II. Pars 2. 8vo.—*From the Editors*.
Abhandlungen der königlich. bayerischen Akademie der
Wissenschaften. Philosophisch-Philologischen Classe—
Band XIII. Abth. 3. Math.-Physik Classe—Band XII.
Abth. 2. Historischen Classe—Band XIII. Abth. 1, 2.
4to.—*From the Academy*.

Munich.—*Catalogus Codicum Manu Scriptorum Bibliothecæ Reciaë Monacensis.* Tomi Primi, Pars Primi, Pars Quarta. 8vo.
—*From the Academy.*

Almanach der königlichen Bayerischen Akademie der Wissenschaften. Jahr 1875. 12mo.—*From the Academy.*

Sitzungsberichte der königl. bayer. Akademie der Wissenschaften. Philosophisch-Philologischen und Historischen Classe 1875—Band II. Heft 1-4; Band I. Heft 1. Mathematisch-Phisikalischen Classe—1875, Heft 2, 3; 1876, Band I. Heft. 8vo.—*From the Academy.*

Neuchatel.—*Bulletin de la Société des Sciences Naturelles de Neuchatel.* Tome X. Part 2. 8vo.—*From the Society.*

New Haven (U.S.)—*Journal (American) of Science and Art,* conducted by Benjamin Silliman. Vol. X. Nos. 58-60; XI. Nos. 61-66; XII. Nos. 67, 68. 8vo.—*From the Editor.*

Transactions of the Connecticut Academy of Arts and Sciences. Vol. III. Part I. 8vo.—*From the Academy.*

New York.—*Fifty-Sixth Annual Report of the Trustees of the New York State Library for the year 1873; Fifty-Seventh Annual Report for the year 1874.* 8vo.—*From the Trustees.*
Eighty-Sixth Annual Report of the Regents of the University for 1873; Eighty-Seventh Annual Report for 1874. 8vo.—*From the University.*

New Zealand.—*Statistics of the Colony of New Zealand, 1874.* Folio.—*From the New Zealand Government.*

Tenth Annual Report of the Colonial Museum and Laboratory. 1874-75. By Dr Hector. 8vo.—*From the Geological Survey.*

Nijmegen.—*Nederlandsch Kruidkundog Archief.* Deel II. Stuk 1. 8vo.—*From the Editors.*

Oxford.—*Astronomical and Meteorological Observations made at the Radcliffe Observatory, Oxford, in the year 1873.* Vol. XXXIII. 8vo.—*From the Observatory.*

Paris.—*Annales des Mines.* Tome VIII. Liv. 4°, 5°, 6°; Tome IX. Liv. 1°. 8vo.—*From the Ecole des Mines.*

Bulletin de la Société de Géographie. 1875, Septembre, Octobre, Novembre, Decembre; 1876, Janvier, Fevrier, Mars, Avril, Mai, Juin. 8vo.—*From the Society.*

Paris.—Bulletin de la Société Mathématique de France. Tome III.

Nos. 3, 5, 6. 8vo.—*From the Society.*

Nouvelles Archives du Museum d'Histoire Naturelle. Tom

X. Fasc. 1-2. 4to.—*From the Natural History Museum.*

Publications of the Dépôt de la Marine (with Charts). 8vo.

—*From the Dépôt.*

Revue Historique Dirigée. Par M.M. G. Monod et G.

Fagniez. Tome Premier, No. 1. 1876. 8vo.—*From the*

Editors.

Revue Philosophique de la France et de l'Étranger Dirigée.

Par Th. Ribot. Première, No. 1. 1876. 8vo.—*From*

the Editor.

Philadelphia.—Proceedings of the American Philosophical Society.

Vol. XIV., Nos. 94 and 95. 8vo.—*From the Society.*

Transactions of the American Philosophical Society. Vol.

XV. New Series, Part 2. 4to.—*From the Society.*

Rome.—Bolletino del Comitato Geologico d'Italia. Nos. 1-6.

8vo.—*From the Society.*

Salem (U.S.)—The American Naturalist. Vols. VIII. and IX.

8vo.—*From the Peabody Academy of Science.*

Memoirs of the American Association for the Advance-

ment of Science. Vol. I. 4to.—*From the Associa-*

tion.

Memoirs of the Peabody Academy of Science. Vol. I.

No. 4. 8vo.—*From the Academy.*

Sixth Annual Report of the Trustees of the Peabody

Academy of Science, for the year 1873. 8vo.—*From the*

Academy.

Sidney.—Proceedings of the Linnean Society of New South Wales.

Vol. I. Part 1. 8vo.—*From the Society.*

Stockholm.—Kongliga Svenska Vetenskaps-Akademienes Hand-

lingar Ny Folja Attönde. Bandet 1870-71-73. 4to.

—*From the Academy.*

Meteorologiska Jakttagelser i Sverige utgifna af Köngl.

Svenska Vetenskaps-Akademiens Anställda och Bear-

betade af Er-Edlund. Tofte Bandet 1870; Trellönde

Bandet 1871; Fjertönde Bandet 1872. 4to.—*From the*

Academy.

Stockholm.—Ofversigt af Köngl. Vetenskaps Akademiens Förhandlingar Tjugondeattonde. Argangen 1871–1872; Tretionde Argangen 1873; Tretiondeforsla Argangen 1874. 8vo.
—*From the Academy.*

Bihang till Köngl. Svenska Vetenskaps-Akademiens Handlingar. Första Bandet, Hafte 1–2; Andra Bandet, Hafte 1–2. 8vo.—*From the Academy.*

Lefnadsteckningar ofver Köngl. Svenska Vetenskaps-Akademienstefter ar 1854, Afidna Ledamoter. Band I. Hefte 3. 8vo.—*From the Academy.*

St Petersburg.—Annalen Physikalischen Central Observatoriums. 1874. 4to.—*From the Russian Government.*

Bulletin de l'Académie Impériale des Sciences de St Petersburg. Tome XXI. No. 5. 4to.—*From the Academy.*

Repertorium für Meteorologie. Band IV. 4to.—*From the Royal Academy.*

Tableau General Methodique et Alphabetique des Matieres Contenues dans les Publications de l'Académie Impériale des Sciences de St Pétersbourg. 1872, 1^{re} Partie. 8vo.
—*From the Academy.*

Toronto.—Canadian Journal of Science, Literature, and History. Vol. XIV. Nos. 5, 6; Vol. XV. Nos. 1, 2. 8vo.—*From the Editor.*

Turin.—Atti della Reale Accademia della Scienze de Torino. Vol. X. Dispensa 1–8. 8vo.—*From the Academy.*

Bolletini dell Osservatorio della Regia Università de Torino. 1875. 4to.—*From the University.*

R. Osservatorio Astronomico di Torino Effemride del Sole, Della Luna e die Principali Pianeti calcolate per Torino, in Tempo Medio Civile di Roma, per l'Anno 1876–1877. 8vo.—*From the University.*

Bolletino Meteorologico ed Astronomico del Regio Osservatorio della Università di Torino. 1873–1874. 4to.—*From the University.*

Upsala.—Nova Acta Regiæ Societatis Scientiarum Upsaliensis. Vol. IX. Fasc. 2. 4to.—*From the Society.*

Bulletin Météorologique Mensuel de l'Observatoire de l'Université d'Upsal. Vol. VI. Année 1874. 4to.—*From the University.*

Utrecht.—Nederlandsch Meteorologisch Jaarboek voor 1870 and 1874. 4to.—*From the Meteorological Institute.*

Verslag van het Verhandelde in de Algemeene Vergadering van het Provinciaal Utrechtsche genootschap van Kunsten en Wetenschappen. 1874. 8vo.—*From the Society of Sciences and Arts.*

Victoria.—Statistical Register of the Colony of Victoria for 1874. Fol. Population—Part 6. Statistical Register Interchange—Part 5. Reports on the Mining Surveyors and Registrars, 1875.—Fol. Patents and Patentees—Vol. VIII.—4to. Statistics of Friendly Societies, 1874.

Statistical Register of the Colony of Victoria.—Part 7. Accumulation. Fol.—*From the Registrar General.*

Vienna.—Das Gebirge um Hallstatt. Eine Geologisch-Paläontologische Studie aus den Alpen. Abhandlungen der k. k. Geologischen Reichsanstalt. Band VI. Heft 2. Edmund Mojsisovics v. Mojsvár. 4to.—*From the Society.*

Die Congerien und Paludinensichten Slavoniens und Eeren Faunen, von D. M. Neumayl und C. M. Paul. Band VII. Heft 3. 4to.—*From the Society.*

Denkschriften der kaiserlichen Academie der Wissenschaften. Math.-Natur. Classe. Band XXXIV. 4to.—*From the Academy.*

Sitzungsberichte der kaiserlichen Academie der Wissenschaften.—Min. Bot Zool. Geo. Pal. Classe. Band LXX. Hefts 3, 4, 5. LXXI. Hefts 1, bis 5.—Phys. Anat. Classe. Band LXX. Hefts 3, bis 5; LXXI. Hefts 1, 2.—Phil. Hist. Classe. LXXVIII. Hefts 2, 3; LXXX. Hefts 1, 2, 3; LXXIX. Hefts 1, 2, 3.—Math. Nat. Classe. LXX. Hefts 3, 4 bis 5; LXXI. Hefts 1, bis 5. 8vo.—*From the Academy.*

Almanach der Kaiserlichen Academie der Wissenschaften. 1875. 8vo.—*From the Academy.*

Jahrbuch der kaiserlich-königlichen Geologischen Reichsanstalt. Band XXV. Nos. 3, 4; XXVI. No. 1. 8vo.—*From the Society.*

Vienna.—Verhandlungen der kaiserlich-königlichen Zoologisch-botanischen Gesellschaft in Wien. Band XXV. 8vo.—
From the Society.

Verhandlungen der kaiserlich-königlichen Geologischen Reichsanstalt. 1875, 1-5; 11, 12, 13. Nos. 14-18. 1876. 8vo.—*From the Society.*

Festschrift zur feier des Funfundzwanzigjährigen Bestehens k. k. Zoologisch-Botanischen Gesellschaft in Wien. 4to.—
From the Society.

Warwick.—Fortieth Annual Report (1876) of the Natural History and Archæological Society. 8vo.—*From the Society.*

Proceedings of the Warwickshire Naturalists' and Archæologists' Field Club. 1875. 8vo.—*From the Society.*

Washington.—Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1874. 8vo.—*From the Institution.*

Astronomical and Meteorological Observations made during the year 1873. 4to.—*From the U. S Naval Observatory.*

Bulletin of the United States Geological and Geographical Survey of the Territories. 1875. Nos. 2, 3. 8vo.—
From the Geological Survey.

Report of the Geographical and Geological Surveys West of the Mississippi. 1875. 8vo.—*From the Geological Survey.*

Report of the United States Geological Survey of the Territories. By F. V. Hayden. Vol. II. 4to.—*From the Geological Survey.*

Annual Report of the United States Geological and Geographical Survey of the Territories, embracing Colorado and parts of adjacent Territories, being a Report of the Exploration for the year 1874. By F. V. Hayden. 8vo.—
From the Geological Survey.

Watford.—Transactions of the Watford Natural History and Hertfordshire Field Club. Vol. I. Part 1-3. 8vo.—*From the Society.*

Wellington (N.Z.)—Reports on the Durability of New Zealand Timber in Constructive Works. 8vo.—*From the Agent General.*

Wellington (N.Z.)—Transactions and Proceedings of the New Zealand Institute. 1875. Vol. VIII. 8vo.—*From the Institute.*

Whitby.—Report of the Literary and Philosophical Society. 1875. 8vo.—*From the Society.*

York.—Annual Report of the Yorkshire Philosophical Society for 1874. 8vo.—*From the Society.*

Zurich.—Schweizerische Meteorologische Beobachtungen. 4to.—*From the Observatory.*

Vierteljahrschrift der Naturforschenden Gesellschaft in Zurich. Neunter Jahrgang, Heft 1-2. 8vo.—*From the Society.*

